

A Proposed



Docket Number	4-31336P1
---------------	-----------

~~jc/59 U.S. PTO~~

FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10

EL402023583US

Express Mail Label Number

March 10, 2000

Date of Deposit

Address to: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

UTILITY PATENT APPLICATION TRANSMITTAL AND FEE SHEET

Transmitted herewith for filing under 37 CFR §1.53(b) is the utility patent application of

Applicant (or identifier): CROWL ET AL.

Title: ARTHRITIS-ASSOCIATED PROTEIN

Enclosed are:

1. ☒ Specification (Including Claims and Abstract) - 32 pages
2. ☒ Drawings - 5 sheets
3. ☒ Executed Declaration and Power of Attorney (original or copy)
4. ☐ Microfiche Computer Program (appendix)
5. ☐ Nucleotide and/or Amino Acid Sequence Submission
 - ☒ Computer Readable Copy
 - ☒ Paper Copy
 - ☒ Statement Verifying Identity of Above Copies
6. ☐ Preliminary Amendment
7. ☐ Assignment Papers (Cover Sheet & Document(s))
8. ☐ English Translation of
9. ☐ Information Disclosure Statement
10. ☐ Certified Copy of Priority Document(s)
11. ☒ Return Receipt Postcard
12. ☐ Other:

Filing fee calculation:

- ☐ Before calculating the filing fee, please enter the enclosed Preliminary Amendment.
☐ Before calculating the filing fee, please cancel claims .

Basic Filing Fee								\$	690
Multiple Dependent Claim Fee (\$ 260)								\$	
Foreign Language Surcharge (\$ 130)								\$	
	For	Number Filed		Number Extra		Rate			
Extra Claims	Total Claims	21	-20	1	x	\$	18	= \$ 18	
	Independent Claims	5	-3	2	x	\$	78	= \$ 156	
TOTAL FILING FEE								\$	864


- ☒ Please charge Deposit Account No. 19-0134 in the name of Novartis Corporation in the amount of \$864. An additional copy of this paper is enclosed. The Commissioner is hereby authorized to charge any additional fees under 37 CFR §1.16 and §1.17 which may be required in connection with this application, or credit any overpayment, to Deposit Account No. 19-0134 in the name of Novartis Corporation.

Please address all correspondence to the address associated with Customer No. 001095, which is currently:

Thomas Hoxie
Novartis Corporation
Patent and Trademark Dept.
564 Morris Avenue
Summit, NJ 07901-1027

Please direct all telephone calls to the undersigned at the number given below, and all telefaxes to (908) 522-6955.

Respectfully submitted,



David E. Wildman
Agent for Applicants
Reg. No. 40,226
Tel. No. (908) 522-6946

Date: March 10, 2000

ARTHRITIS-ASSOCIATED PROTEIN

FIELD OF THE INVENTION

This invention relates to methods and compositions for early diagnosis, monitoring and treatment of cartilage degenerative conditions, including forms of arthritis. In particular, the invention relates to a protein differentially expressed in human subjects suffering from arthritis, such as osteoarthritis (OA) and rheumatoid arthritis (RA) compared to individuals without arthritis, antibodies that recognize this protein, and methods for diagnosing arthritis.

BACKGROUND OF THE INVENTION

Arthritis involves both cartilage breakdown and new bone formation, and in many cases, leads to the loss of joint function. While there are some insights as to the mediators contributing to cartilage remodeling, there is only limited knowledge of the underlying molecular mechanisms of arthritis. At a cellular level, the articular chondrocyte plays a key role in both cartilage breakdown and new bone growth, and this is likely to be reflected in changes in cellular transcriptional activity. The application of genomic approaches to the understanding of the complexities of arthritis promises to yield significant advances in the diagnosis and treatment of this prevalent disease of joint degeneration.

An important priority for the investigation and clinical application of potential therapies for arthritis is the identification of biochemical markers that could be used to assess various aspects of disease activity (Chevalier, X. (1997) *Rev.Rhum. Engl. Ed.* **64**: 562-577.

Lohmander, L.S. (1997) *Baillieres Clin. Rheumatol.* **11**: 711-726). Current research in this field is focusing on a number of possible surrogate markers of arthritis that reflect metabolic changes in the joint associated with cartilage destruction and remodeling, including hyaluronate, cartilage oligomeric matrix protein, keratan sulfate, metalloproteinase activity, and various cytokines.

To identify gene products involved in human arthritis, the present inventors have analyzed the differences in mRNA populations in diseased cartilage compared to age-matched healthy cartilage using differential display RT-PCR (Hu, S-I, Carozza, M., Klein, M., Nantermet, P., Luk, D., and Crawl, R.M. (1998) *J. Biol. Chem.* **273**: 34406-34412). The inventors have

discovered a large protein, termed "adlican," the presence of which in cartilage is strongly associated with arthritis, particularly osteoarthritis. An anti-peptide antibody was generated against the carboxy terminal sequence and used for immunoblot analysis to detect adlican protein in synovial fluid samples obtained from patients with arthritis.

SUMMARY OF THE INVENTION

In a first aspect, the invention provides an isolated DNA comprising a nucleotide sequence as set forth in SEQ ID NO:1. Also provided are isolated DNA's comprising nucleic acid sequences that hybridizes under high stringency conditions to the isolated DNA as set forth in SEQ ID NO:1. In a preferred embodiment, the isolated DNA takes the form of a vector molecule comprising the DNA as set forth in SEQ ID NO:1.

In a second aspect, the invention provides an isolated polypeptide with an amino acid sequence as set forth in SEQ ID NO:2. Such a polypeptide, or fragments thereof, is found in the synovial fluid of sufferers of arthritis to a much greater extent than in the synovial fluid of individuals without arthritis. In accordance with this aspect of the invention there are provided novel polypeptides of human origin as well as biologically, diagnostically or therapeutically useful fragments, variants and derivatives thereof, variants and derivatives of the fragments, and analogs of the foregoing.

A third aspect of the present invention encompasses a method for the diagnosis of arthritis in a human which includes detecting the elevated transcription of messenger RNA transcribed from adlican-encoding DNA in cartilage from a human, where such elevated transcription is diagnostic of the human's affliction with arthritis.

Another embodiment of the assay aspect of the invention provides a method for the diagnosis of arthritis in a human which requires measuring the amount of a polypeptide that includes or is adlican or fragments of adlican in synovial fluid from a human, where the presence of an elevated amount of the polypeptide or fragments thereof, relative to the amount of the polypeptide or fragments thereof in non-arthritic synovial fluid, is diagnostic of the human's suffering from arthritis.

In accordance with one embodiment of this aspect of the invention there is provided anti-sense polynucleotides that regulate transcription of the human adlican gene.

It is another object of the invention to provide a process for producing the
aforementioned polypeptides, polypeptide fragments, variants and derivatives, fragments of the
variants and derivatives, and analogs of the foregoing. In a preferred embodiment of this aspect
of the invention there are provided methods for producing the aforementioned human adlican
polypeptides comprising culturing host cells having incorporated therein an expression vector
containing an exogenously-derived human adlican-encoding polynucleotide under conditions
sufficient for expression of human adlican polypeptides in the host and then recovering the
expressed polypeptide.

In accordance with another object the invention there are provided products,
compositions, processes and methods that utilize the aforementioned polypeptides and
polynucleotides for research, biological, clinical and therapeutic purposes, inter alia.

In certain additional preferred embodiments of this aspect of the invention there are
provided an antibody or a fragment thereof which specifically binds to a polypeptide that
comprises the amino acid sequence set forth in SEQ ID NO:2, i.e., adlican. In certain
particularly preferred embodiments in this regard, the antibodies are highly selective for human
adlican polypeptides or portions of human adlican polypeptides.

In a further aspect, an antibody or fragment thereof is provided that binds to a fragment
or portion of the amino acid sequence set forth in SEQ ID NO:2.

In yet another aspect, the invention provides vertebrate cells which can be propagated in
vitro and which are capable upon growth in culture of producing a polypeptide that comprises
the amino acid sequence set forth in SEQ ID NO:2 or fragments thereof, where the cells contain
transcriptional control DNA sequences, other than human adlican transcriptional control
sequences, where the transcriptional control sequences control transcription of DNA encoding a
polypeptide with the amino acid sequence according to SEQ ID NO: 2 or fragments thereof.

In another aspect, the present invention provides a method for producing human adlican
polypeptides which comprises culturing a host cell having incorporated therein an expression
vector containing an exogenously-derived human adlican-encoding polynucleotide under
conditions sufficient for expression of human adlican polypeptides in the host cell, thereby
causing the production of an expressed polypeptide, and recovering the expressed polypeptide.

In yet another aspect of the present invention there are provided kits comprising the components necessary to detect above-normal expression of human adlcan polynucleotides or polypeptides in body tissue samples derived from a patient.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a depiction of the cDNA encoding adlcan (SEQ ID NO:1).

Figure 2 is a depiction of the amino acid sequence encoded by SEQ ID NO:1 (SEQ ID NO:2).

Figure 3 is an analysis of the amount of adlcan mRNA expressed in the cartilage of human osteoarthritis sufferers compared with the amount of adlcan mRNA expressed in the cartilage of human non-arthritic control subjects using semi-quantitative reverse transcription-PCR. Semi-quantitative reverse-transcription PCR of actin was used as a control.

Figure 4 is an autoradiogram of a gel showing the production of adlcan *in vitro* using a reticulocyte lysate system.

Figure 5 is a Western Blot of a gel on which proteins from HEK 293 cells which were transfected with an adlcan expression vector were run. Anti-adlcan peptide antibodies were used. Lane 1 is cell extract from cells transfected with an adlcan expression vector 72 hours post-infection. Lane 2 is from cells 48 hours post-infection. Lane 3 is from cells transfected with a control vector at 48 hours post-infection.

Figure 6 is a Western Blot of a gel on which synovial fluid samples from humans suffering from trauma, osteoarthritis, and rheumatoid arthritis were tested. Anti-adlcan peptide antibodies were used.

DETAILED DESCRIPTION OF THE INVENTION

All patent applications, patents and literature references cited herein are hereby incorporated by reference in their entirety.

In practicing the present invention, many conventional techniques in molecular biology, microbiology, and recombinant DNA are used. These techniques are well known and are explained in, for example, Current Protocols in Molecular Biology, Volumes I, II, and III, 1997 (F. M. Ausubel ed.); Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; DNA Cloning: A Practical Approach, Volumes I and II, 1985 (D. N. Glover ed.); Oligonucleotide Synthesis, 1984 (M. L. Gait ed.); Nucleic Acid Hybridization, 1985, (Hames and Higgins); Transcription and Translation, 1984 (Hames and Higgins eds.); Animal Cell Culture, 1986 (R. I. Freshney ed.); Immobilized Cells and Enzymes, 1986 (IRL Press); Perbal, 1984, A Practical Guide to Molecular Cloning; the series, Methods in Enzymology (Academic Press, Inc.); Gene Transfer Vectors for Mammalian Cells, 1987 (J. H. Miller and M. P. Calos eds., Cold Spring Harbor Laboratory); and Methods in Enzymology Vol. 154 and Vol. 155 (Wu and Grossman, and Wu, eds., respectively).

As used herein, "differentially expressed gene" refers to (a) a gene containing at least one of the DNA sequences disclosed herein (e.g., as shown in Figure 1 and SEQ ID NO:1); (b) any DNA sequence that encodes the amino acid sequence encoded by the DNA sequences disclosed herein (e.g., as shown in Figure 2 and SEQ ID NO:2); or (c) any DNA sequence that is substantially similar to the coding sequences disclosed herein.

In its broadest sense, the term "substantially similar", when used herein with respect to a nucleotide sequence, means a nucleotide sequence corresponding to a reference nucleotide sequence, wherein the corresponding sequence encodes a polypeptide having substantially the same structure and function as the polypeptide encoded by the reference nucleotide sequence, e.g. where only changes in amino acids not affecting the polypeptide function occur. Desirably the substantially similar nucleotide sequence encodes the polypeptide encoded by the reference nucleotide sequence. The percentage of identity between the substantially similar nucleotide sequence and the reference nucleotide sequence desirably is at least 80%, more desirably at least 85%, preferably at least 90%, more preferably at least 95%, still more preferably at least 99%.

Sequence comparisons are carried out using a Smith-Waterman sequence alignment algorithm (see e.g. Waterman, M.S. Introduction to Computational Biology: Maps, sequences and genomes. Chapman & Hall. London: 1995. ISBN 0-412-99391-0, or at <http://www-hto.usc.edu/software/seqaln/index.html>).

The localS program, version 1.16, is used with following parameters: match: 1, mismatch penalty: 0.33, open-gap penalty: 2, extended-gap penalty: 2. A nucleotide sequence "substantially similar" to reference nucleotide sequence hybridizes to the reference nucleotide sequence in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 2X SSC, 0.1% SDS at 50°C, more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 1X SSC, 0.1% SDS at 50°C, more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.5X SSC, 0.1% SDS at 50°C, preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 50°C, more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50°C with washing in 0.1X SSC, 0.1% SDS at 65°C, yet still encodes a functionally equivalent gene product.

The differentially expressed genes disclosed herein are expressed in arthritic tissues (e.g., cartilage in a human afflicted with arthritis) in elevated amounts relative to, i.e., to a greater extent than in the corresponding tissues of humans who do not suffer from arthritis. Messenger RNA transcribed from the differentially expressed genes, and protein translated from such mRNA, is present in arthritic tissues and/or synovial fluid associated with such tissues in an amount at least about twice, preferably at least about five times, more preferably at least amount ten times, most preferably at least about 100 times the levels of mRNA and protein found in corresponding tissues found in humans who do not suffer from arthritis. Such elevated transcription of adlcan mRNA is referred to herein as "elevated transcription."

A "host cell," as used herein, refers to a prokaryotic or eukaryotic cell that contains heterologous DNA that has been introduced into the cell by any means, e.g., electroporation, calcium phosphate precipitation, microinjection, transformation, viral infection, and the like.

"Heterologous" as used herein means "of different natural origin" or represent a non-natural state. For example, if a host cell is transformed with a DNA or gene derived from another organism, particularly from another species, that gene is heterologous with respect to

that host cell and also with respect to descendants of the host cell which carry that gene. Similarly, heterologous refers to a nucleotide sequence derived from and inserted into the same natural, original cell type, but which is present in a non-natural state, e.g. a different copy number, or under the control of different regulatory elements.

5 A vector molecule is a DNA molecule into which heterologous DNA may be inserted which can then replicate in an appropriate host cell. Vectors must have one or more origin of replication, and one or more site into which the recombinant DNA can be inserted. Vectors often have convenient means by which cells with vectors can be selected from those without, e.g., they encode drug resistance genes. Common vectors include plasmids, viral genomes, and (primarily
10 in yeast) "artificial chromosomes."

"Plasmids" generally are designated herein by a lower case p preceded and/or followed by capital letters and/or numbers, in accordance with standard naming conventions that are familiar to those of skill in the art. Starting plasmids disclosed herein are either commercially available, publicly available on an unrestricted basis, or can be constructed from available
15 plasmids by routine application of well known, published procedures. Many plasmids and other cloning and expression vectors that can be used in accordance with the present invention are well known and readily available to those of skill in the art. Moreover, those of skill readily may construct any number of other plasmids suitable for use in the invention. The properties, construction and use of such plasmids, as well as other vectors, in the present invention will be
20 readily apparent to those of skill from the present disclosure.

The term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the
25 natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

As used herein, the term "transcriptional control sequence" refers to DNA sequences, such as initiator sequences, enhancer sequences, and promoter sequences, which induce, repress,

or otherwise control the transcription of protein encoding nucleic acid sequences to which they are operably linked.

As used herein, "human adlcan transcriptional control sequences" are any of those transcriptional control sequences normally found associated with the human adlcan gene in as it is found in the human X chromosome.

As used herein, "non-human transcriptional control sequence" is any transcriptional control sequence not found in the human genome.

The term "polypeptide" is used interchangeably herein with the terms "polypeptides" and "protein(s)".

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (c), in the preceding paragraph. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6.times.SSC/0.05% sodium pyrophosphate at 37 °C. (for 14-base oligos), 48 °C. (for 17-base oligos), 55 °C. (for 20-base oligos), and 60 °C. (for 23-base oligos). These nucleic acid molecules may act as target gene antisense molecules, useful, for example, in target gene regulation and/or as antisense primers in amplification reactions of target gene nucleic acid sequences. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for target gene regulation. Still further, such molecules may be used as components of diagnostic methods whereby the presence of a cardiovascular disease-causing allele, may be detected.

The invention also encompasses (a) DNA vectors that contain any of the foregoing coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and (c) genetically engineered host cells that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression.

The invention includes fragments of any of the DNA sequences disclosed herein. Fragments of the full length adlcan gene may be used as a hybridization probe for a cDNA library to isolate the full length gene and to isolate other genes which have a high sequence similarity to the adlcan gene or similar biological activity. Probes of this type preferably have at least about 30 bases and may contain, for example, from about 30 to about 50 bases, about 50 to about 100 bases, about 100 to about 200 bases, or more than 200 bases. The probe may also be used to identify a cDNA clone corresponding to a full length transcript and a genomic clone or clones that contain the complete ADLican gene including regulatory and promotor regions, exons, and introns. An example of a screen comprises isolating the coding region of the ADLican gene by using the known DNA sequence to synthesize an oligonucleotide probe. Labeled oligonucleotides having a sequence complementary to that of the gene of the present invention are used to screen a library of human cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

In addition to the gene sequences described above, homologs of such sequences, as may, for example be present in other species, may be identified and may be readily isolated, without undue experimentation, by molecular biological techniques well known in the art. Further, there may exist genes at other genetic loci within the genome that encode proteins which have extensive homology to one or more domains of such gene products. These genes may also be identified via similar techniques.

For example, the isolated differentially expressed gene sequence may be labeled and used to screen a cDNA library constructed from mRNA obtained from the organism of interest. Hybridization conditions will be of a lower stringency when the cDNA library was derived from an organism different from the type of organism from which the labeled sequence was derived. Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Such low stringency conditions will be well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Further, a previously unknown differentially expressed gene-type sequence may be isolated by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of amino acid sequences within the gene of interest. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express a differentially expressed gene allele.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of a differentially expressed gene-like nucleic acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled fragment may be used to screen a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source. A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies which may be used, see e.g., Sambrook et al., 1989, supra.

In cases where the differentially expressed gene identified is the normal, or wild type, gene, this gene may be used to isolate mutant alleles of the gene. Such an isolation is preferable in processes and disorders which are known or suspected to have a genetic basis. Mutant alleles may be isolated from individuals either known or suspected to have a genotype which contributes to cardiovascular disease symptoms. Mutant alleles and mutant allele products may then be utilized in the diagnostic assay systems described below.

A cDNA of the mutant gene may be isolated, for example, by using PCR, a technique which is well known to those of skill in the art. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying the mutant allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized

using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant gene to that of the normal gene, the mutation(s) responsible for the loss or alteration of function of the mutant gene product can be ascertained.

Alternatively, a genomic or cDNA library can be constructed and screened using DNA or RNA, respectively, from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. The normal gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant allele in the library. The clone containing this gene may then be purified through methods routinely practiced in the art, and subjected to sequence analysis as described above.

Additionally, an expression library can be constructed utilizing DNA isolated from or cDNA synthesized from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal gene product, as described, below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.) In cases where the mutation results in an expressed gene product with altered function (e.g., as a result of a missense mutation), a polyclonal set of antibodies are likely to cross-react with the mutant gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis as described above.

Differentially expressed gene products include those proteins encoded by nucleotide sequence set forth in SEQ ID NO: 1, in particular, a polypeptide that is or includes the amino acid sequence set out in SEQ ID NO: 2, or fragments thereof.

In addition, differentially expressed gene products may include proteins that represent functionally equivalent gene products. Such an equivalent differentially expressed gene product may contain deletions, additions or substitutions of amino acid residues within the amino acid sequence encoded by the differentially expressed gene sequences described, above, but which result in a silent change, thus producing a functionally equivalent differentially expressed on

pathway gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved.

For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Functionally equivalent", as utilized herein, refers to a protein capable of exhibiting a substantially similar *in vivo* activity as the endogenous differentially expressed gene products encoded by the differentially expressed gene sequences described above. Alternatively, when utilized as part of assays such as those described, below, "functionally equivalent" may refer to peptides capable of interacting with other cellular or extracellular molecules in a manner substantially similar to the way in which the corresponding portion of the endogenous differentially expressed gene product would.

The differentially expressed gene products may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the differentially expressed gene polypeptides and peptides of the invention by expressing nucleic acid encoding differentially expressed gene sequences are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing differentially expressed gene protein coding sequences and appropriate transcriptional/translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, *supra*, and Ausubel et al., 1989, *supra*. Alternatively, RNA capable of encoding differentially expressed gene protein sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, M. J. ed., IRL Press, Oxford, which is incorporated by reference herein in its entirety.

A variety of host-expression vector systems may be utilized to express the differentially expressed gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified,

but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the differentially expressed gene protein of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing differentially expressed gene protein coding sequences; yeast (e.g. *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing the differentially expressed gene protein coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the differentially expressed gene protein coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing differentially expressed gene protein coding sequences; or mammalian cell systems (e.g. COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothioneine promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

Expression of adlcan by a cell from an adlcan gene that is native to a the cell can also be performed. Methods for such expression are detailed in, e.g., U.S. Patents 5,641,670; 5,733,761; 5,968,502; and 5,994,127, all of which are expressly incorporated by reference herein in their entirety.

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the differentially expressed gene protein being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of antibodies or to screen peptide libraries, for example, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the differentially expressed gene protein coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. PGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general,

such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene protein can be released from the GST moiety.

Promoter regions can be selected from any desired gene using vectors that contain a reporter transcription unit lacking a promoter region, such as a chloramphenicol acetyl transferase ("cat") transcription unit, downstream of restriction site or sites for introducing a candidate promoter fragment; i.e., a fragment that may contain a promoter. As is well known, introduction into the vector of a promoter-containing fragment at the restriction site upstream of the cat gene engenders production of CAT activity, which can be detected by standard CAT assays. Vectors suitable to this end are well known and readily available. Two such vectors are pKK232-8 and pCM7. Thus, promoters for expression of polynucleotides of the present invention include not only well known and readily available promoters, but also promoters that readily may be obtained by the foregoing technique, using a reporter gene.

Among known bacterial promoters suitable for expression of polynucleotides and polypeptides in accordance with the present invention are the E. coli lacI and lacZ promoters, the T3 and T7 promoters, the T5 tac promoter, the lambda PR, PL promoters and the trp promoter. Among known eukaryotic promoters suitable in this regard are the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus ("RSV"), and metallothionein promoters, such as the mouse metallothionein-I promoter.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is one of several insect systems that can be used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The differentially expressed gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of differentially expressed gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used

to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Pat. No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the differentially expressed gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing differentially expressed gene protein in infected hosts. (E.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted differentially expressed gene coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire differentially expressed gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the differentially expressed gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., 1987, Methods in Enzymol. 153:516-544).

Selection of appropriate vectors and promoters for expression in a host cell is a well known procedure and the requisite techniques for expression vector construction, introduction of the vector into the host and expression in the host per se are routine skills in the art.

Generally, recombinant expression vectors will include origins of replication, a promoter derived from a highly-expressed gene to direct transcription of a downstream structural sequence, and a selectable marker to permit isolation of vector containing cells after exposure to the vector.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, etc.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the differentially expressed gene protein may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the differentially expressed gene protein. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed gene protein.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk^- , $hgprt^-$ or $aprt^-$ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for $dhfr$, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt , which confers

resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hygro, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147) genes.

5 An alternative fusion protein system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88: 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus
10 are loaded onto Ni^{2+} nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

When used as a component in assay systems such as those described below, the differentially expressed gene protein may be labeled, either directly or indirectly, to facilitate detection of a complex formed between the differentially expressed gene protein and a test
15 substance. Any of a variety of suitable labeling systems may be used including but not limited to radioisotopes such as ^{125}I ; enzyme labeling systems that generate a detectable calorimetric signal or light when exposed to substrate; and fluorescent labels.

Where recombinant DNA technology is used to produce the differentially expressed gene protein for such assay systems, it may be advantageous to engineer fusion proteins that can
20 facilitate labeling, immobilization and/or detection.

Indirect labeling involves the use of a protein, such as a labeled antibody, which specifically binds to either a differentially expressed gene product. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by an Fab expression library.

25 Described herein are methods for the production of antibodies capable of specifically recognizing one or more differentially expressed gene epitopes. Such antibodies may include, but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')_2 fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any
30 of the above. Such antibodies may be used, for example, in the detection of a fingerprint, target,

gene in a biological sample, or, alternatively, as a method for the inhibition of abnormal target gene activity. Thus, such antibodies may be utilized as part of cardiovascular disease treatment methods, and/or may be used as part of diagnostic techniques whereby patients may be tested for abnormal levels of adlcan, or for the presence of abnormal forms of adlcan.

For the production of antibodies to a differentially expressed gene, various host animals may be immunized by injection with a differentially expressed gene protein, or a portion thereof. Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as target gene product, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with differentially expressed gene product supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to the hybridoma technique of Kohler and Milstein, (1975, *Nature* 256:495-497; and U.S. Pat. No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, *Immunology Today* 4:72; Cole et al., 1983, *Proc. Natl. Acad. Sci. USA* 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, *Proc. Natl. Acad. Sci.*, 81:6851-6855; Neuberger et al., 1984, *Nature*, 312:604-608;

Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) can be adapted to produce differentially expressed gene-single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Particularly preferred, for ease of detection, is the sandwich assay, of which a number of variations exist, all of which are intended to be encompassed by the present invention.

For example, in a typical forward assay, unlabeled antibody is immobilized on a solid substrate and the sample to be tested brought into contact with the bound molecule. After a suitable period of incubation, for a period of time sufficient to allow formation of an antibody-antigen binary complex. At this point, a second antibody, labeled with a reporter molecule capable of inducing a detectable signal, is then added and incubated, allowing time sufficient for the formation of a ternary complex of antibody-antigen-labeled antibody. Any unreacted material is washed away, and the presence of the antigen is determined by observation of a signal, or may be quantitated by comparing with a control sample containing known amounts of antigen. Variations on the forward assay include the simultaneous assay, in which both sample and antibody are added simultaneously to the bound antibody, or a reverse assay in which the labeled antibody and sample to be tested are first combined, incubated and added to the unlabeled

surface bound antibody. These techniques are well known to those skilled in the art, and the possibility of minor variations will be readily apparent. As used herein, "sandwich assay" is intended to encompass all variations on the basic two-site technique. For the immunoassays of the present invention, the only limiting factor is that the labeled antibody be an adlcan-specific antibody.

The most commonly used reporter molecules in this type of assay are either enzymes, fluorophore- or radionuclide-containing molecules. In the case of an enzyme immunoassay an enzyme is conjugated to the second antibody, usually by means of glutaraldehyde or periodate. As will be readily recognized, however, a wide variety of different ligation techniques exist, which are well-known to the skilled artisan. Commonly used enzymes include horseradish peroxidase, glucose oxidase, beta-galactosidase and alkaline phosphatase, among others. The substrates to be used with the specific enzymes are generally chosen for the production, upon hydrolysis by the corresponding enzyme, of a detectable color change. For example, p-nitrophenyl phosphate is suitable for use with alkaline phosphatase conjugates; for peroxidase conjugates, 1,2-phenylenediamine or toluidine are commonly used. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates noted above. A solution containing the appropriate substrate is then added to the tertiary complex of antibody-adlcan-labeled antibody. The substrate reacts with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an evaluation of the amount of adlcan which is present in the serum sample.

Alternately, fluorescent compounds, such as fluorescein and rhodamine, may be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody absorbs the light energy, inducing a state of excitability in the molecule, followed by emission of the light at a characteristic longer wavelength. The emission appears as a characteristic color visually detectable with a light microscope. Immunofluorescence and EIA techniques are both very well established in the art and are particularly preferred for the present method. However, other reporter molecules, such as radioisotopes, chemiluminescent or bioluminescent molecules may

also be employed. It will be readily apparent to the skilled artisan how to vary the procedure to suit the required use.

The present invention also includes a kit for performing the assay aspect of the invention. Such a kit includes vials or vessels for incubating a body tissue or fluid sample, and the components necessary for quantifying human adlcan polynucleotides, for example, via RT-PCR. A kit for quantifying human adlcan polypeptide may contain anti-adlcan antibodies, for example, the antibodies may be prepared via the procedure set forth in Example 5.

The following Examples illustrate the present invention, without in any way limiting the scope thereof.

EXAMPLES

Example 1: Cloning of Adlcan

Differential display RT-PCR is used to identify gene products with altered expression in diseased cartilage from humans suffering from OA compared to cartilage from non-arthritic (NA) human controls. As described by Hu *et al.*, above, first strand cDNA is synthesized from 0.2 µg of total RNA with each of the 3 anchored oligo-dT primers from GenHunter Corporation, Nashville, TN. The reaction (20 µl) was carried out at 37 °C for 60 min. For PCR amplification, 1 µl of the cDNA serves as template in a 10 µl reaction mix containing 10 mM Tris-HCl (pH 8.4), 1.5 mM MgCl₂, 50 mM KCl, 0.001% gelatin, 2 µM dNTPs, 0.2 µM of 5' arbitrary primer (AP-3 from the RNAimage kit obtained from GenHunter Corporation, Nashville, TN), 2 µM of the same anchored primer used in the cDNA synthesis, 5 µCi of α-[³³P]dATP (2,000 Ci/mmol, Dupont-New England Nuclear, Boston, MA) and 2.5 units of AmpliTaq DNA polymerase (Perkin-Elmer, Norwalk, CT). Samples are subjected to 40 cycles of amplification under the following conditions: denaturing at 94 °C, 30 sec, annealing at 40 °C, 2 min, extension at 72 °C, 30 sec, and a final extension at 72 °C, 5 min. The resulting PCR products are resolved on a denaturing polyacrylamide gel and visualized by autoradiography of the dried gel. PCR products of interest are excised from the gel, and the DNA is eluted and re-amplified by PCR using the same primers and conditions described above, excluding the radio-labeled nucleotide. PCR products are analyzed on a 1.5% agarose gel and are ligated into the cloning vector PCR II 2.1

(TA cloning Kit, Invitrogen, Carlsbad, CA). Clones of the PCR-generated fragments are obtained by transformation of *E. coli* strain DH5 α (Gibco/BRL, Gaithersburg, MD). DNA sequences are determined for at least 3 independent clones of each fragment using Dye Terminator Cycle Sequencing on an ABI PRISM 377 DNA sequencing system (Perkin-Elmer, Norwalk, CT).

As described by Hu *et al*, above, first strand cDNA is synthesized from total RNA isolated from OA and non-arthritic cartilage. 200 ng of total RNA and 10 pmoles of primer T₃₀VN (where V=A,C,G and N=A,C,G,T) are mixed in a 6 μ l volume, heated to 72°C for 3 min and quenched on ice for 3 minutes. Buffer and MMLV reverse transcriptase are added to final concentrations of 50 mM Tris-HCl (pH 8.3), 6 mM MgCl₂, 75 mM KCl, 1 mM dNTPs, and 10 units MMLV reverse transcriptase in 10 μ l. This mixture is incubated at 42 °C for 1.5 hr, 94 °C for 5 min, and quenched on ice. Serial dilutions of cDNA from different individuals are used for PCR amplification with a primer set for actin (forward: GGAGTCCTGTGGCATCCACGAAACTAC (SEQ ID NO:3) and reverse: CACATCTGCTGGAAGGTGGACAGCG (SEQ ID NO:4) and for 3G12 (forward: CATGGGCACAACTGACTCATGGCTG (SEQ ID NO:5) and reverse: GAGAGGAGAGGAAGGAGAAAAGTGCATC (SEQ ID NO:6)) under the following conditions: 25 μ l reaction volume with 10 mM Tris-HCl (pH 8.3) , 50 mM KCl, 1.5 mM MgCl₂, 0.001% gelatin, 20 μ M of dNTPs, 1.25 units AmpliTaq Gold Polymerase (Perkin Elmer, Norwalk, CT); 94 °C, 8.5 minutes; 32 cycles of 94 °C, 30 seconds; 63 °C, 30 seconds; 72 °C, 2 minutes, and a final incubation at 72 °C, 7 minutes. Nine μ l of the reaction mix is run on a 10% polyacrylamide gel, stained with SYBR™ Green I (Molecular Probes, Eugene, OR), and quantified using fluorescence imaging and ImageQuant software (Molecular Dynamics, Sunnyvale, CA). The resulting PCR fragment for 3G12 is 155 bp. Concentrations are chosen to ensure that the reaction is within the log phase of amplification. Levels of expression are normalized to the levels of actin for each sample. A negative control reaction with no template is carried out for each primer set to verify the absence of contamination.

From this screening approach, several differentially expressed transcripts are obtained, including a 251 bp clone designated 3G12 (detected with 5' primer AP-3 and 3' primer oligo dT with G, 12th clone isolated). The sequence of 3G12 is determined to be a novel sequence identified in the screen. Using PCR primers designed based on the 3G12 sequence, the

corresponding transcript is found to be expressed over 100 times greater in OA cartilage than in NA cartilage by semi-quantitative RT-PCR (Figure 3). Because of the high degree of differential expression, 3G12 is pursued as a diagnostic marker gene for OA.

The cloned 251 bp PCR-generated 3G12 fragment is used as a probe to identify a λ gt11 phage clone designated "106A" with a 1.4 kb insert in a human placenta cDNA library. From a human placental 5'-STRETCH PLUS cDNA Library (Oligo dT + random primed, Clontech, Palo Alto, CA), 10^7 pfu were divided into 40 sub-libraries and screened by PCR using 3G12-specific primers. Positive fractions were further divided into 21 second-round sub-libraries which were also screened by PCR. Phage from the positive second-round sub-libraries were plated on *E. coli* strain Y1090 and screened by plaque hybridisation using a PCR-generated probe (251 bp) labeled with [32 P] dATP. Filters were hybridized at 38 °C for 2.5 hours and washed at 48 °C three times in 1x SSC containing 0.1% SDS. Positive plaques were identified by autoradiography, isolated and amplified, and re-screened until positive single phage clones were confirmed.

A BLASTX analysis of the sequence of 106A identifies a partial open reading frame (ORF) with coding sequence related to the L1 neural cell adhesion molecule (Coutelle, O., Nyakatura, G., Taudien, S., Elgar, G., Brenner, S., Platzer, M., Drescher, B., Jouet, M., Kenwick, S., and Rosenthal, A. (1998) *Gene* **208**: 7-15), neurofascin (Hassel, B., Rathjen, F.G., and Volkmer, H. (1997) *J Biol Chem* **272**: 28742-28749), and to perlecan, a large heparan sulfate containing proteoglycan (Murdoch, A.D., Dodge, G.R., Cohen, I., Tuan, R.S., Iozzo, R.V. (1992) *J Biol Chem* **267**: 8544-8557). Although it is evident that 106A represents a partial transcript, no cDNA clones are yet identified with additional upstream sequences. A repeated search of the Genbank and the EST sequence databases is conducted, and a small number of EST sequences corresponding to 3G12 are found. However, none of these sequences include the 5'-segment of cDNA clone 106A.

Clone 106A is used to identify a human genomic clone (Genome Systems, St. Louis, MO, P1 clone # 14414). Sub-clones of the P1 isolate are obtained, are characterized by DNA sequence analysis, and are found to contain 2 complete and 1 incomplete exons. The 5'-most exon region is then used as a probe to identify another cDNA clone from the placental library,

designated N3. The sequence of N3 extends the ORF, revealing additional homologies to NCAM and perlecan; however, the sequence lacks any evident translational initiation site.

An approach known as 5' RACE (rapid amplification of cDNA ends) is then used to obtain extended clones. Separate first strand cDNA syntheses are carried out with 500 ng of total RNA isolated and pooled from 4 OA cartilage samples. Aliquots are combined for the synthesis of double stranded cDNA with anchors ligated at both ends according to the Marathon™ cDNA system (Clontech, Palo Alto, CA). The antisense primer is derived from the sequence of cDNA clone N3. Utilization of the 5' RACE method yields an additional ~500 bp of coding sequence, although still with no evident translational start site. The 5'-most sequence from the RACE clones is used to isolate a second genomic clone, and its sub-clone E3. A search of Genbank reveals that the sequence of E3 is 100% identical to a 3 kb segment of a sequence deposited into Genbank (Accession number AC004616,), human genomic clone BAC536K7 (149,752 bp) containing a segment of the X chromosome.

The sequence of BAC536K7 is extensively analysed for predicted exons encoding the transcript corresponding to clone 3G12. Various exon prediction programs (e.g. GENESCAN, GRAIL) are used to find the rest of the gene encoding 3G12. PCR experiments are required to test for the presence of putative exons using the original placental cDNA library as template. The 5' exon (X1, exons are referred to 5' to 3' in their order in the gene, X1, X2, etc.) predicted by GENESCAN to contain the translation start site is found to be incorrect. Only by visual inspection using ORF FINDER of small potential coding exons, which contain apparent secretory signal sequences connected to sequences homologous to proteins in common with a down stream exon is the correct 5' exon identified. Exons X1, X2, X3, and X4 are verified by PCR using human placental cDNA as template and primers which bridged pairs of exons. The complete sequence of the 3G12 transcript is derived from 6 exons (X5 and X6 represented by clones N3 and 106A). The entire DNA sequence and encoded protein sequence are shown in Figures 1 and 2, respectively. The large adlcan protein (2,828 amino acids) contains five leucine-rich repeats in the amino terminal region downstream of the signal sequence, distinct regions of homology to neural cell adhesion molecule and perlecan containing immunoglobulin C2 domains, and a mucin-like domain. Based on its predicted functionalities, the protein is

termed “adlican” – for *AD*hesion protein with *Le*ucine-rich repeats and *Im*munoglobulin domains related to perle~~CAN~~.

Example 2: Assembly of Expression Construct for Adlican

In order to produce adlican protein and examine its biochemical properties, multiple sub-cloning steps are required to assemble the various cDNA and genomic clone fragments, comprising the entire coding region, into an expression vector. A first construct is made as follows. A λ gt11 clone which contains 347 bp of 5' UTR and 1-715 bp of coding region is used as a template for PCR. The PCR primer sequences used were:

Upstream (contains a NotI site and the initiation codon ATG)

5'- TTGCGGCCGCGCCACCATGCCCAAGCGCGCGCACTGG-3' (SEQ ID NO:7);

and

Downstream

5'-TCAATACTCCTGCTCCTGTTCTGTCTCA-3' (SEQ ID NO:8).

During the PCR reaction a NotI site is added at the 5'-end to enable subcloning into an expression vector and the 3'-end contained an EcoRI site which is present in the original adlican sequence. The 715bp NotI-EcoRI fragment is subcloned into the pcDNA 3 vector (Invitrogen, Carlsbad, CA) and the nucleotide sequence is verified using an Applied Biosystems 373 automated DNA sequencer and dideoxy terminator chemistry. This fragment also contains a signal sequence of 19 amino acids.

The following cDNA clones are used to generate a second construct. Clone E3 which contains coding regions from 1513-5482 bp is cut with HindIII and EcoRI and a 3969 bp fragment which corresponds to base pairs 1513-5162 of adlican is subcloned into the pcDNA3 vector. A HindIII-HindIII fragment which contains sequences corresponding to base pairs 734-1512bp (778bp) of adlican is generated by PCR using the D5 clone as a template, subcloned and sequenced. The vector that contains the 3969 base pair EcoRI- HindIII fragment is cut with HindIII and the 778 bp HindIII-HindIII fragment is ligated into it. The clone containing the HindIII fragment in the correct orientation contains cDNA sequence corresponding to base pairs

734-5162 of the original sequence. A 19 base pair EcoRI – HindIII adaptor which contains sequence corresponding to base pairs 716-733 is synthesized and ligated to construct II which is partially digested with HindIII to open up the 5' HindIII site located at bp734. This completes construction of the second construct, which contains an insert of 4446 bp (base pairs 716-5162) of the original sequence.

The third construct is generated from E3 and two other cDNA clones N3 and 106A. E3 is cut with EcoRI-HindIII to generate a 320 bp fragment corresponding to base pairs 5163-5482 of adlcan. N3 is cut with HindIII-ApaLI to generate a 297 bp fragment corresponding to base pairs 5483-5780 of adlcan. Clone 106A is cut with ApaLI- XhoI to generate a 2707 bp fragment which corresponds to base pairs 5780-8487 of adlcan. This fragment also contains a stop codon at base pair 8484. The EcoRI- HindIII, HindIII- ApaLI and ApaLI-XhoI fragments are put together by a trimolecular ligation and are subcloned into EcoRI- XhoI cut pcDNA3 to generate a 3325 bp fragment, i.e., the third construct, corresponding to base pairs 5162-8487 of adlcan.

In order to generate the full length clone, the third construct is cut with EcoRI and NotI and the NotI-EcoRI insert of the first construct from above is ligated into it to generate a fourth construct of 4040 bp. The second construct from above is cut with EcoRI and the 4446 bp insert is gel purified.

The fourth construct is digested with EcoRI and the 4446 bp EcoRI insert of the second construct is ligated into it to generate the final full length fifth construct. Clones which have the EcoRI fragment of the second construct in the right orientation represent the full length clone for adlcan. The stop codon is removed from the full length insert and further subcloned in frame into a hexa-histidine tag containing vector (Invitrogen, Carlsbad, CA) for *in vitro* and *in vivo* expression.

Example 3: Expression of the full-length construct in a TNT reticulocyte lysate coupled Transcription/Translation system

The plasmid DNA containing the full length insert is translated in a TNT® coupled reticulocyte lysate system (Promega, Madison, WI) as described by the manufacturer. The reaction mixture containing TNT® rabbit reticulocyte lysate, TNT® reaction buffer, T7 RNA polymerase, amino acid mixture minus methionine, [³⁵S]- methionine (> 1000Ci/mmol) Rnasin-

ribonuclease inhibitor, 0.5µg of pcDNA 3.1 (Invitrogen, Carlsbad, CA), DNA containing the full length adlcan insert and transcend biotin-lysyl-tRNA is incubated at 30 °C for 60-90 minutes. At the end of the incubation an aliquot of the reaction is denatured in SDS containing sample buffer and electrophoresed on a 4-20% gradient polyacrylamide gel. The gel is dried after
 5 soaking for 5 minutes in 7% acetic acid, 7% methanol, 1% glycerol containing buffer and translated products are visualized by autoradiography.

Example 4: Transient expression of the full-length construct in HEK 293 cells

HEK 293 cells are seeded (200,000/well) into 6 well plates in DMEM medium containing
 10 10% FBS and are incubated at 37 °C for 48-72 hours until the cells are about 70% confluent. The cells are transfected with the full length adlcan construct in the pcDNA 3.1 vector using lipofectamine/Plus reagent (Gibco/BRL, Gaithersburg, MD). The DNA is precomplexed with the Plus reagent for 15 minutes and then is incubated with diluted lipofectamine reagent for a further 15 minutes. The cells are washed and transferred to serum-free medium and the
 5 DNA/lipofectamine mixture is added and incubated for 3 hours at 37 °C. The transfection medium is removed and is replaced with complete medium (DMEM containing 10% FBS). The cells are incubated overnight. The cells are shifted into DMEM containing 0.2% FBS the next day and are incubated for a further 48 hours. Aliquots of the supernatant are collected 48 hours and 72 hours post-transfection, are denatured and are electrophoresed on a 4-20% gradient
 20 polyacrylamide gel. The proteins in the gel were electroblotted on to a PVDF membrane (BioRad, Hercules, CA) and probed with anti hexa-his antibody (Qiagen, Valencia, CA). The bound antibody was visualized using the ECL reagent system (Amersham, Piscataway, NJ).

Example 5: Generation of Antibodies that Recognize and Bind to Adlcan

25 An anti-peptide antibody is generated against a sequence near the carboxy-terminus of adlcan based on the predicted coding sequence of the first cDNA clone, 106A. A 15 residue peptide (CMAKNILGSDSKTTY), corresponding to the sequence of the adlcan protein near the carboxyl terminus, is designed based on surface probability as determined using the program PROTEAN, a component of the LASERGENE suite of programs (DNASTAR Inc., Madison,
 30 WI). The peptide is synthesized, purified, and used to immunize two rabbits following a standard

protocol (Genosys Biotechnologies, The Woodlands, TX). Antiserum from one of the immunized animals which shows the highest titer against the peptide is used for immunoblot analysis. Samples (4 μ l) of human synovial fluid are subjected to SDS polyacrylamide gel electrophoresis, are transferred to PVDF membranes (BioRad, Hercules, CA), are analyzed for immunoreactivity using a 1:500 dilution of the antiserum, using the ECL detection system (chemiluminescent) (Amersham, Piscataway, NJ).

The resulting antiserum is used for detecting adlcan protein in human synovial fluid by immunoblot analysis. The data in Fig. 6 include an example of one of the immunoblots where adlcan protein fragment is detected in human synovial fluid and a tabulation of results derived from several blots where synovial fluids from individual patients with varied diagnoses are tested. The diagnoses are osteoarthritis, rheumatoid arthritis, gout, and joint trauma. 100% of the patients with severe OA show a positive adlcan protein immunoreactivity.

What is claimed is:

1. An isolated DNA comprising a nucleotide sequence as set forth in SEQ ID NO:1.
2. A host cell comprising an isolated DNA according to claim 1.
3. A vector molecule comprising at least a fragment of an isolated DNA according to claim 1.
4. A vector molecule according to claim 3 comprising transcriptional control sequences.
5. An isolated polypeptide comprising the amino acid sequence set forth in SEQ ID NO:2.
6. An isolated polypeptide consisting of the amino acid sequence set forth in SEQ ID NO:2.
7. An isolated DNA comprising a nucleic acid sequence that encodes the polypeptide of claim 6.
8. An isolated DNA comprising a nucleic acid sequence of from about 30 to about 50 nucleotides that hybridizes under high stringency conditions to the isolated DNA of claim 1.
9. A host cell comprising a vector molecule according to claim 3.
10. A vertebrate host cell which can be propagated in vitro and which is capable upon growth in culture of producing a polypeptide according to claim 5, wherein said

cell comprises at least one transcriptional control sequence that is not a human adlcan transcriptional control sequence, wherein said one or more transcriptional control sequences control transcription of DNA encoding a polypeptide according to claim 5.

5 11. A vertebrate cell according to claim 10 wherein said one or more transcriptional control DNA sequences are non-human transcriptional control sequences.

 12. A method for the diagnosis of arthritis in a human which comprises:
 detecting the elevated transcription of messenger RNA transcribed from
10 the DNA of claim 7 in cartilage from a human, wherein said elevated transcription is diagnostic of said human's suffering from arthritis.

 13. The method of claim 12, wherein said DNA has the nucleotide sequence set forth in SEQ ID NO:1.

 14. A method for the diagnosis of arthritis in a human which comprises:
 measuring the amount of a polypeptide that comprises the polypeptide
 according to claim 5, or fragments thereof, in synovial fluid from a human, wherein the presence
 of an elevated amount of said polypeptide or fragments thereof, relative to the amount of said
 polypeptide or fragments thereof in non-arthritis synovial fluid, is diagnostic of said human's
 suffering from arthritis.

 15. The method of claim 14, wherein said detecting step comprises contacting
 said synovial fluid with an antibody which specifically binds to a polypeptide that comprises the
25 amino acid sequence set forth in SEQ ID NO:2 or a fragment thereof and detecting specific
 binding of said antibody with a polypeptide in said synovial fluid, wherein detection of specific
 binding to a polypeptide indicates the presence of a polypeptide that comprises the amino acid
 sequence set forth in SEQ ID NO:2 or a fragment thereof.

16. A purified antibody or a fragment thereof which specifically binds to a polypeptide that comprises the amino acid sequence set forth in SEQ ID NO:2 or a fragment of a polypeptide that comprises the amino acid sequence set forth in SEQ ID NO:2.

5 17. An antibody fragment according to claim 16 which is an Fab or F(ab')₂ fragment.

18. An antibody according to claim 16 which is a polyclonal antibody.

10 19. An antibody according to claim 16 which is a monoclonal antibody.

20. A method for producing human adlcan polypeptides which comprises:
culturing a host cell having incorporated therein an expression vector containing
an exogenously-derived human adlcan-encoding polynucleotide under conditions sufficient for
expression of human adlcan polypeptides in the host cell, thereby causing the production of an
expressed polypeptide; and
recovering the polypeptide produced by said cell.

21. An isolated DNA molecule with a nucleotide sequence complementary to
the nucleotide sequence of the isolated DNA according to claim 1.

ABSTRACT

Disclosed are methods and compositions for early diagnosis, monitoring and treatment of cartilage degenerative conditions, including forms of arthritis. In particular, the invention relates to a protein, termed "adlican," that is differentially transcribed and expressed in human subjects suffering from arthritis, such as osteoarthritis (OA) and rheumatoid arthritis (RA) compared with non-arthritis sufferers, antibodies which recognize this protein, and methods for diagnosing arthritis.

Patent 6,636,336

FIG. 1

ATGCCCAAGCGCGCGCACTGGGGGGCCCTCTCCGTGGTGCTGATCCTGCTTTGGGGCCATCCGCGAGTGGCGCTGGCCTGCCCGCATCCTTGTGCCT
GCTACGTCCCCAGCGAGGTCCACTGCACGTTCCGATCCCTGGCTTCCGTGCCCGCTGGCATTGCTAGACACGTGGAAGAATCAATTTGGGGTTAA
TAGATACAGGCCCTGTACAGAAACCTCATTGGCAGGACTGACCAAGTTGGAGCTACTATTGATTCCCGCAATGAGATCCCAAGCATCCCGATGGA
GCTTTAAGAGACCTCAGCTCTCTTCAGGTTTTCAAGTTCAGCTACAACAAGCTGAGAGTGATCACAGGACAGACCCCTCCAGGCTCTCTCTAACTTAA
TGAGGCTGCACATTGACCACAACAAGATCGAGTTTATCCACCCTCAAGCTTTCAACGGCTTAACGCTCTCTGAGGCTACTCCATTTGGAAGGAAATCT
CCTCCACCAGCTGCACCCAGCACCTTCTCCACGTTACATTTTTGGATTATTTAGACTCTCCACCATAAGGCACCTCTACTTAGCAGAGAACATG
GTTAGAATCTTCCCTGCCAGCATGCTTCGGAACATGCCGCTTCTGGAGAATCTTTACTTGCAGGGAAATCCGTGGACCTGCGATTGTGAGATGAGAT
GGTTTTTGGAAATGGGATGCAAAATCCAGAGGAATTTGAAGTGTAAGAAAGGACAAAGCTTATGAAGGCGGTCAAGTTGTGTCAATGTGCTTCAGTCC
AAAGAAGTTGTACAAACATGAGATACACAAGCTGAAGGACATGACTTGTCTGAAGCCTTCAATAGAGTCCCTCTGAGACAGAACAGGAGCAGGAGT
ATTGAGGAGGAGCAAGAACAGGAAGAGGATGGTGGCAGCCAGCTCATCTGGAGAATTTCCAACCTGCCCCAGTGGAGCATCTCTTGAATATGACCG
ACGAGCACGGGAACATGGTGAACCTGGTCTGTGACATCAAGAAACCAATGGATGTGTACAAAGATTCACTTGAACCAACCGGATCCCTGAGATGAGAT
CATAAATGCAACAGTTGCCTTGGACTTTGAGTGTCCAATGACCCGAGAAAATATGAAAAGCTATGGAAATGATAGCATACTACAGTGAAGTTCCC
GTGAGCTACACAGAGAGTCTCATCTCAGCAAAAGACCCAGAGTCAGTACCAGTACAGGAGGATGCTGATGAGGAAGCTCTTTACTACACAGGTG
TGAGAGCCAGATTCTTGCAACCAAGATGGGTCTGACAGCCATCCATAGATATCCAGCTGAACCGACGTCAGAGTACGGCCAAAGAGGTGCTACT
TTCTACTACACCTGTTTCTCAAACATATCCACCAAGATAGACAGGCTCGGGGAGAGCTGGGTAATGATTAGGCTTAGTGGAGCTGTG
CAAAGAGATCAGACTGTCTTGGAAAGGGGTCCATGCCAGTTGACGTGCAACGCTTCTGAGAGTCCATCTATCTTCTGGGTGCTTCCAGATG
GCTCCATCTGAAAGCGCCCATGGATGACCCAGACAGCAAGTTCTCCATTCTCAGCAGTGGCTGGCTGAGGATCAAGTCCATGGAGCCATCTGACTC
AGGCTTGTACCAAGTGCATTGCTCAAGTGAAGGATGAATGGACCGCATGGTATATAGGGTACTTGTGCACTCTCCCTCCACTCAGCCAGCCGAGAAA
ACAGAGGTGAATTAATGATTTGGCTAAACACATCAGATGTATACATGTGCAAAAGTGAAGTCTTTCCATCCCAAGGTCAAGTCAAGTATGAGTGG
TTACTACAGATGTGTGGCTGTCAACCAGCAAGGGGAGACCATTTTACGGTGGGAATCACAGTGACCAAGAAAGGGTCTGGCTTGGCATCCAAAAGA
GGCAGACGCCAGGTGCAAGGGTCTTTCCAGAGTCAGAGAAGACATCGTGGAGGATGAAGGGGGCTCGGGCATGGGAGATGAAGAGAACACTTCAA
GGAGACTTCTGCAATCCAAAGGACCAAGAGGTGTCTCTCAAAGAGGATGATGCCATCAATGGAGACAAGAAAGGAGGAGAGAAAGCT
GAAACTCTGGAAGCATTCGGAAGAAAGAACAGAGACCAATGTTGCAAGAGTGCAGAGTGTGATCTGAAATCTAGACGAGGATAAATGGAAGCAACAA
CAGATTAATCCGGAGCGTGGGCTGATATTTAGCCAAAGTCCGTGGGAAAATCTCCCTAAGGGCAGAGAAGTACCCCGTTGATTAAGAACACAA
GTCCTCCATCTTGAAGCTAGAGTCAACACACCTTTTCTGCTGTTTCTCCCCCTCAGCATCTCTGTGCAGACAGTAAACAGTGTGAAGATC
CTCAGCAGATGTACCTTACTTGGTGAAGAAGACAGCTTTTGGGTACCATTCTCAGCCAGCATGGGGCTAGAACACAACCAATGGAGTTATT
CTTGTGAAACCTGAAGTAAACAAGCACACCTCTGGAGGAAGTTGTGAGTACCTTCTGAGAGACTGAGGAGATAACTTCCACTGAAGGAGACCTGA
AGGGGACAGCAGCCCTACACTTATATCTGAGCCTTATGAACCATCTCTACTCTGCACACATTAGACACAGTCTATGAAAGGCCACCCATGAAGA
GACGGCAACAGAGGGTGTGCTGCAGCAGATGTGGATCGTCAACAGAGCCACATCCAGTGAGTATGAGCTCCATTGGATGCTGTCTCTTGGCT
GAGTCTGAGCCCATGCAATACTTTGACCCAGATTGGAGACTAAGTCAACAGGATGAGGATAAGATGAAGAAGACACCTTTGCACACCTTACTC
CAACCCCCACCATCTGGGTAACTGACTCCAGTACATCAGTATTATTGAGGATTTACTATAGGGGAACAGGTGTCCAGGCCATCTACATACA
AGGACTGACAGACAACATCCACCTTGTGAAAGTAGTCTAAGCACTCAAGACACCTTACTGATTAAGAGGGTATGAAGAGATGTCTCAGACACTA
CAGGAGGAGAAATGTCTAGAGGAGAGCCACACACTCCAGAGTCTGAGAGTGAGGGCAAGAGAGCAAAATCCATCACTTTGGCTGACTCCACAC
TGGGTATAATGAGCAGTATGTCTCCAGTTAAGAAGCCTGCGGAAACACAGTTGGTACCCTCTAGACAAAGACACCACAACAGTAACAACACACC
AAGGCAAAAGTTGCTCCGTCACTCCACCATCAGCCTCACCCTCTGAGAGGAGACCAACGGGAGAGAGATTACGAGGATTTCCGCTTCCGCTC
CGGCACAAGCAACCCACCCACAACCTTTGCCCCATCAGAGACTTTTCTACTCAACCAACTCAAGCACCTGACATTAAGATTTCAAGTCAAGTGG
AGATTTCTCTGGTTCTACAGCTTGGGTGGATAACACAGTTAATACCCCAACAGTTGAAATGGAGAAGATGCAAGAACCCATCCAGGGGAAC
ACCAGGAGAAAACACGGGAAGAGGCAAAACACATCGATATACCCCTTCTACAGTGAGCTCAAGAGCGTCCGGATCCAAGCCAGCCCTTCTCCA
GAAATAAACATAGAAACATTTGTTACTCCAGTTCAGAAACTATATCTTGGCTGAAACTGAGGCTGTTTCTGAAAACTGAGGCGCTTATGTTCTTAG
ATTACATGACAACCACAGAAAATATATTATCTTACCCTAAGTCCAAGAGACACTTCCAGTCACATATAAACCACATCAGATGGAAGAGAAAT
TAAGGATGATGTTGCCACAATGTTGACAACATAAAGTGACATTTTAGTCACTGGTGAATCAATTAATGCTATACCAACTTCTCGCTCCTTG
GTCTCCACTATGGGAGATTTAAGGAAGAACTCTCTCTGAGGCTTCCAGGAATCCAACTGGAATCCCTCAAGGACGGCCAGCCTGGGAGGC
TACAGACAGACATACCTGTTACCACTTCTGGGGAAAATCTTACAGACCTCCCTTCTTAAAGAGCTTGAGGATTTCCCTTCCGATTTCTT
GTCTCTTTGACAGTCTCCACACCATTTACCAGGAAGAAGCTGGTCTTCCACAACCTCTCTCAAGCATAAAGTGGAGTGGCTTCAAGTCAGGCA
GAAACCACACCCCTTGATCAAGATCATCTTGAACCACTGTGGCTATTCTCTTCTGAAACTAGACCACAGAATCACACCCCTACTGCTGCCCGGA
TGAAGGAGCCAGCATCTCTGCTCCCATCCACAATTCTCATGTCTTTGGGACAACACCACCACCTAAGCCAGCACTTCCAGTCCAAGATATCTCA
AGCATCTAGAGATTCCAAGGAAATGTTTCTTGAATTAATGTTGGGAATCCAGAAACAGCAACCCCACTCAACAATGAAGGAACACAGCATATG
TCAGGGCCAAATGAATTATCAACACCTCTTCCGACCGGATGCATTTAAGTGTCTACAAAGCTGGAATGGAAAAGCAAGTATTGTTAGTAGGA
GTCTACCAGTGGCCAGATAGCCAACGCCAGGATGGAAGAGTTCATGCTTCTCATCAACTAACCAGAGTCCCTGCCAAACCCATCTTACCAACAGC
ACAGTGAGGCTACCTGAAATGTCACACAAGCGCTTCCAGATACCTTGTAACTTCCAGTCACCTCGTCACTGGACCAACAACCCGGAATAACT
ACATATCCTTCTGGGGCTTGGCAGAGAACAAACAGTTTACAACCTCAAGATTATCAAGTACAACAATTCCTCTCCCATTTGCACATGTCCAAACCCA
GCATTCCTAGTAAGTTTACTGACCGAAGAACTGACCAATTCAATGGTTACTCCTCAAGTGTGTTGGAATAACAACATCCCTGAGGCAAGAAACCCAGT
TGGAAAGCCTCCAGTCCAAGAAATCTCTATTATCCAATGGAAGACTCCCTTCTTACCAACAAGACTCTTCTTTTCCACAGTTGGGAGTCACC
CGGAGACCCAGATACCCACTTCTCTGCCCCAGTAAAGAGAGAGAAAAGTTATCCAGGTTCTTACACAGGATACATTCATAGCACCTTCC
ATCTGGACTTTGGCCCTCCGGCACTCCGCTTGTGACACTCCGAGACACGGGATCACCCTCAACTAAGTTACAGAAATCTCTTATGTTCTCTTCC
CACCCAGAGTTCTATCTCTTTATAACATCTTCTGTCCAGTCTCAGGAAGCTTCCACCAGAGCAGCTCAAAGTTCTTTGAGGAGGACCTCTGCA
TCCAATTTCTGGTCTTCTGGGGAAGAGCCCAATCTCACCAGTCCCCACAGACTGTGTCCGTACCCGTGAGACAGACAGTGTGTTCCCTGTG
AGGCAACAGGAAAACCAAGCCTTCTGTTACTTGGACAAGGTTTCCACAGAGCTCTTATGACTCCGAATACAGGATACAAAGGTTTGGGTTCT
CAAGAACAGTACTTAGTGATACGGAAGGTTCAAGTACAAGTTCAGAGACTGAGAAACATCTGTCACCGCCAGCAACCTGACAGGCTGAGTG
GTCTTGTCTTCTGGTCAACGTGCAGCAACCTCAATCTAGCTCCCACTACCAGGAGCTCACTGTCTACCTGGGAGACACCATGCAATGGAGTGTCT
TGGCCAAAGGGACCCAGCCCCCAATTTCTGATCTTCCCTGACAGGAGGGTGTGGCAACTGTGTCCCGGTGGAGAGCCGATCACCCTGCA
CGAAACCGGACCCCTTCCATCAAGGAGGCTCTCTCAGACAGAGGCTCTATAAGTGGTGGCCAGCAATGCAGCCGGGGGCGAGCCTGGCC
ATCCGCTGCACGTGGCGGCACTGCCCGGTTATCCACAGGAGAGCTGAGAAACATCTCGCTGCCCGGGGCTCAGACTTCACTTCACTGCA
CTGCCAAGGCTGCGCCCTGCCAGCGTGCCTGGGTGCTCGGGACGGTACCAGATCCGCCCTCGAGTCTCTCCACGGGAACCTGTTTGTGTTT
CCCCAACGGGACGCTTACATCCGCAACCTCGCGCCCAAGGACAGCGGGCGTATGAGTGCCTGGCCGCCAACCTGGTAGGCTCCGCGCGCAGGACG
GTGCAGCTGAACGTGCAGCGTGCAGCAGCAACCGCGCATCACGGGACCTCCCGCGGAGGACGAGCTCAGGTACGGAGGAACCTCAAGCTGG

FIG. 3




FIG. 4




FIG. 5





FIG. 6



DECLARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATIONS

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,
and

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled

ARTHRITIS-ASSOCIATED PROTEIN

the specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims.

I acknowledge my duty to disclose all information which is known by me to be material to the patentability of this application as defined in 37 C.F.R. §1.56.

I hereby claim the benefit under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate listed below and under 35 U.S.C. §365(a) of any PCT international application(s) designating at least one country other than the United States listed below and have also listed below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application the priority of which is claimed for that subject matter:

None

I hereby claim the benefit under 35 USC §119(e) of any United States provisional application(s) listed below:

None

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) listed below and under 35 U.S.C. §365(c) of any PCT international application(s) designating the United States listed below and, insofar as the subject matter of each of the claims of this

application is not disclosed in said prior application(s) in the manner required by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose all information known by me to be material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date(s) of the prior application(s) and the national or PCT international filing date of this application:


None

I hereby appoint the attorneys and agents associated with Customer No. 001095, respectively and individually, as my attorneys and agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.


Please address all communications to the address associated with Customer No. 001095, which is currently Thomas Hoxie, Novartis Corporation, Patent and Trademark Dept., 564 Morris Avenue, Summit, NJ 07901-1027.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FIRST JOINT INVENTOR:

Full name : Robert Mitchell Crowl
Signature : 
Date : 03/09/00
(MM/DD/YY)
Citizenship : United States of America
Residence : Berkeley Heights, New Jersey
P.O. Address : 16 Wentworth Drive
Berkeley Heights, New Jersey 07922

SECOND JOINT INVENTOR:

Full name : Daniel C. Luk
Signature : 
Date : 03/09/2000
(MM/DD/YY)
Citizenship : United States of America
Residence : Passaic, New Jersey
P.O. Address : 5 Barry Place
Passaic, New Jersey 07055

IMPORTANT: Before this declaration is signed, the patent application (the specification, the claims and this declaration) must be read and understood by each person signing it, and no changes may be made in the application after this declaration has been signed.

Novartis Corporation
Patent and Trademark Dept.
564 Morris Avenue
Summit, NJ 07901-1027
(908) 522-6946
Date: March 10, 2000

<110> Crawl, Robert M.
Luk, Daniel C.

<130> 4-31336

<160> 8

<170> FastSEO for Windows Version 4.0

<210> 1

<211> 9645

<212> DNA

<213> Human

<400> 1

atgcccaagc	gcgcgcactg	gggggcccctc	tccgtggtgc	tgatcctgct	ttggggccat	60
ccgcgagtg	gcgtggcctg	cccgcatcct	tgtgctgtgc	acgtccccag	cgaggtccac	120
tgcacgttcc	gatccctggc	ttccgtgccc	gctggcattg	ctagacacgt	ggaaagaatc	180
aatttgggg	ttaatagcat	acaggccctg	tcagaaacct	catttgcagg	actgaccaag	240
ttggagctac	ttatgattca	cggcaatgag	atcccaagca	tccccgatgg	agctttaaga	300
gacctcagct	ctcttcagg	tttcaagttc	agctacaaca	agctgagagt	gatcacagga	360
cagaccctcc	agggtctctc	taacttaatg	aggctgcaca	ttgaccacaa	caagatcgag	420
tttatccacc	ctcaagcttt	caacggctta	acgtctctga	ggctactcca	tttgggaagga	480
aatctcctcc	accagctgca	ccccagcacc	ttctccacgt	tcacattttt	ggattattttc	540
agactctcca	ccataaggca	cctctactta	gcagagaaca	tggttagaac	tcttcctgcc	600
agcatgcttc	ggaacatgcc	gcttctggag	aatctttact	tgcagggaaa	tccgtggacc	660
tgcgattgtg	agatgagatg	gtttttggaa	tgggatgcaa	aatccagagg	aattctgaag	720
tgtaaaaagg	acaaagctta	tgaaggcggt	cagttgtgtg	caatgtgctt	cagtccaaag	780
aagttgtaca	aacatgagat	acacaagctg	aaggacatga	cttgtctgaa	gccttcaata	840
gagtcccctc	tgagacagaa	caggagcagg	agtattgagg	aggagcaaga	acaggaagag	900
gatggtggca	gccagctcat	cctggagaaa	ttccaactgc	cccagtggag	catctctttg	960
aatatgaccg	acgagcaagg	gaacatggtg	aacttgggtc	gtgacatcaa	gaaaccaatg	1020
gatgtgtaca	agattcactt	gaaccaaacy	gatcctccag	atattgacat	aaatgcaaca	1080
gttgcttgg	actttgagtg	tccaatgacc	cgagaaaact	atgaaaagct	atggaaattg	1140
atagcatact	acagtgaagt	tcccgatgaag	ctacacagag	agctcatgct	cagcaaagac	1200
cccagagtca	gctaccagta	caggcaggat	gctgatgagg	aagctcttta	ctacacaggt	1260
gtgagagccc	agattcttgc	agaaccagaa	tgggtcatgc	agccatccat	agatatccag	1320
ctgaaccgac	gtcagagtac	ggccaagaag	gtgctacttt	cctactacac	ccagtattct	1380
caaacaatat	ccaccaaaaga	tacaaggcag	gctcggggca	gaagctgggt	aatgattgag	1440
cctagtggag	ctgtgcaaaag	agatcagact	gtcctggaag	ggggtccatg	ccagttgagc	1500
tgcaacgtga	aagcttctga	gagtccatct	atcttctggg	tgcttccaga	tggtccatc	1560
ctgaaagcgc	ccatggatga	cccagacagc	aagttctcca	ttctcagcag	tggtggctg	1620
aggatcaagt	ccatggagcc	atctgactca	ggcttgatcc	agtgcattgc	tcaagtgagg	1680
gatgaaatgg	accgcatgg	atatagggtc	cttggtgcagt	ctccctccac	tcagccagcc	1740
gagaaagaca	cagtgacaat	tggcaagaac	ccaggggagt	cggtgacatt	gccttgcaat	1800
gctttagcaa	taccgaagc	ccaccttagc	tggattcttc	caaacagaag	gataattaat	1860
gatttggcta	acacatcaca	tgtatacatg	ttgccaaatg	gaactctttc	catcccaaag	1920

gtccaagtca	gtgatagtgg	ttactacaga	tgtgtggctg	tcaaccagca	aggggcagac	1980
cattttacgg	tgggaatcac	agtgaccaag	aaagggctctg	gcttgccatc	caaaagaggc	2040
agacgcccg	gtgcaaaggc	tctttccaga	gtcagagaag	acatcgtgga	ggatgaaggg	2100
ggctcgggca	tgggagatga	agagaacact	tcaaggagac	ttctgcatcc	aaaggaccaa	2160
gaggtgttcc	tcaaaacaaa	ggatgatgcc	atcaatggag	acaagaaagc	caagaaaggg	2220
agaagaaagc	tgaaactctg	gaagcattcg	gaaaaagaac	cagagaccaa	tgttgcagaa	2280
ggtcgcagag	tgtttgaatc	tagacgaagg	ataaacatgg	caaacaaaca	gattaatccg	2340
gagcgtctgg	ctgatatattt	agccaaagtc	cgtgggaaaa	atctccctaa	gggcacagaa	2400
gtacccccgt	tgattaaaac	cacaagtcct	ccatccttga	gcctagaagt	cacaccacct	2460
tttcctgctg	tttctccccc	ctcagcatct	cctgtgcaga	cagtaaccag	tgtctgaagaa	2520
tcctcagcag	atgtacctct	acttggtgaa	gaagagcacg	ttttgggtac	catttcctca	2580
gccagcatgg	ggctagaaca	caaccacaat	ggagttattc	ttgttgaaac	tgaagtaaca	2640
agcacacctc	tggaggaagt	tgttgatgac	ctttctgaga	agactgagga	gataacttcc	2700
actgaaggag	acctgaaggg	gacagcagcc	cctacactta	tatctgagcc	ttatgaacca	2760
tctcctactc	tgcacacatt	agacacagtc	tatgaaaagc	ccacccatga	agagacggca	2820
acagaggggt	ggctctgcagc	agatgttgga	togtcaccag	agcccacatc	cagtgagtat	2880
gagcctccat	tggatgctgt	ctccttggct	gagtcctgagc	ccatgcaata	ctttgaccca	2940
gatttggaga	ctaagtcaca	accagatgag	gataagatga	aagaagacac	ctttgcacac	3000
cttactccaa	ccccaccat	ctgggttaat	gactccagta	catcacagtt	atttgaggat	3060
tctactatag	gggaaccagg	tgtcccaggc	caatcacatc	tacaaggact	gacagacaa	3120
atccaccttg	tgaaaagtag	tctaagcact	caagacacct	tactgattaa	aaagggtatg	3180
aaagagatgt	ctcagacact	acagggagga	aatatgctag	agggagaccc	cacacactcc	3240
agaagtctctg	agagtggagg	ccaagagagc	aaatccatca	ctttgcctga	ctccacactg	3300
ggtataatga	gcagtatgtc	tccagttaag	aagcctgogg	aaaccacagt	tggtagccctc	3360
ctagacaaaag	acaccacaac	agtaacaaca	acaccaaggc	aaaaagttgc	tccgtcatcc	3420
accatgagca	ctcacccttc	tccaaggaga	cccaacggga	gaaggagatt	acgccccaac	3480
aaattccggc	accggcacia	gcaaacccca	cccacaactt	ttgccccatc	agagactttt	3540
tctactcaac	caactcaagc	acctgacatt	aagattttcaa	gtcaagtggga	gagttctctg	3600
gttctctacag	cttgggtgga	taacacagtt	aataccccca	aacagttgga	aatggagaag	3660
aatgcagaac	ccacatccaa	gggaacacca	cggagaaaaac	acgggaagag	gccaaacaaa	3720
catcgatata	ccccttctac	agtgagctca	agagcgtccg	gatccaagcc	cagcccttct	3780
ccagaaaata	aacatagaaa	cattgtttact	cccagttcag	aaactatact	tttgcttaga	3840
actgtttctc	tgaaaactga	gggcccttat	gattccttag	attacatgac	aaccaccaga	3900
aaaatatatt	catcttacct	taaagtccaa	gagacacttc	cagtcacata	taaaccaca	3960
tcagatggaa	aagaaattaa	ggatgatgtt	gccacaaatg	ttgacaaaaca	taaaagtgc	4020
atthtagtca	ctggtgaatc	aattactaat	gccataccaa	cttctcgtct	cttgggtctcc	4080
actatgggag	aatttaagga	agaatcctct	cctgtaggct	ttccaggaac	tccaaccttg	4140
aatccctcaa	ggacggccca	gcctgggagg	ctacagacag	acatacctgt	taccacttct	4200
ggggaaaaatc	ttacagaccc	tccccttctt	aaagagcttg	aggatgtgga	tttacttcc	4260
gagtttttgt	cctctttgac	agtctccaca	ccatttcacc	aggaagaagc	tggttcttcc	4320
acaactctct	caagcataaa	agtggaggtg	gcttcaagtc	aggcagaaac	cggaccctt	4380
gatcaagatc	atcttgaaac	cactgtggct	attctccttt	ctgaaactag	accacagaat	4440
cacaccctta	ctgctgcccg	catgaaggag	ccagcatcct	cgtccccatc	cacaattctc	4500
atgtcttttg	gacaaaccac	gaccactaag	ccagcacttc	ccagtccaag	aatatctcaa	4560
gcactctagag	attccaagga	aaatgttttc	ttgaattatg	tggggaatcc	agaaacagaa	4620
gcaaccccag	ttaacaatga	aggaacacag	catatgtcag	ggccaaatga	attatcaaca	4680
ccctcttccg	accgggatgc	atttaacttg	tctacaaaagc	tgggaattgga	aaagcaagta	4740
tttggtagta	ggagtctacc	acgtggccca	gatagccaac	gccaggatgg	aagagttcat	4800
gcttctcatc	aactaaccag	agtccctgcc	aaacccatcc	taccaacagc	aacagtgagg	4860
ctacctgaaa	tgtccacaca	aagcgcttcc	agatactttg	taacttccca	gtcactcgt	4920
cactggacca	acaaaaccgga	aataactaca	tatccttctg	gggctttgcc	agagaacaaa	4980
cagttttacaa	ctccaagatt	atcaagtaca	acaattctct	tcccattgca	catgtccaaa	5040
cccaqcattc	ctagtaagtt	tactgaccga	agaactgacc	aattcaatgg	ttactccaaa	5100

gtgttttgga	ataacaacat	ccctgaggca	agaaaccag	ttggaagcc	tcccagtcca	5160
agaattcctc	attattccaa	tggaagactc	cctttcttta	ccaacaagac	tctttctttt	5220
ccacagttgg	gagtcacccg	gagaccccag	ataccactt	ctcctgccc	agtaatgaga	5280
gagagaaaag	ttattccagg	ttcctacaac	aggatacatt	cccatagcac	cttccatctg	5340
gactttggcc	ctccggcacc	tccgttggtg	cacactccgc	agaccacggg	atcacccctca	5400
actaacttac	agaatatccc	tatgggtctct	tccaccaga	gttctatctc	ctttataaca	5460
tcttctgtcc	agtcctcagg	aagcttccac	cagagcagct	caaagttctt	tgcaggagga	5520
cctcctgcat	ccaaattctg	gtctcttggg	gaaaagcccc	aaatcctcac	caagtcccca	5580
cagactgtgt	ccgtcacccg	tgagacagac	actgtgttcc	cctgtgaggc	aacaggaaaa	5640
ccaaagcctt	tcgttacttg	gacaaagggt	tccacaggag	ctcttatgac	tccgaatacc	5700
aggatacaac	ggtttgagggt	tctcaagaac	ggtaccttag	tgatacggaa	ggttcaagta	5760
caagatcgag	gccagtatat	gtgcaccgcc	agcaacctgc	acggcctgga	caggatgggtg	5820
gtcttgcttt	cggtcacccg	gcagcaacct	caaatcctag	cctcccacta	ccaggacgtc	5880
actgtctacc	tgggagacac	cattgcaatg	gagtgtctgg	ccaaagggac	cccagccccc	5940
caaatttcct	ggatcttccc	tgacaggagg	gtgtggcaaa	ctgtgtcccc	cgtggagagc	6000
cgcatacccc	tgcacgaaaa	ccggaccctt	tccatcaagg	aggcgctcct	ctcagacaga	6060
ggcgtctata	agtgcgtggc	cagcaatgca	gccggggcgg	acagcctggc	catccgcctg	6120
cacgtggcgg	cactgcccc	cgttatccac	caggagaagc	tggagaacat	ctcgtgccc	6180
ccggggctca	gcattcacat	tactgcact	gccaaaggctg	cgccccctgc	cagcgtgccc	6240
tgggtgctcg	gggacggtag	ccagatccgc	ccctcgcagt	tctccacgg	gaacttgttt	6300
gttttcccca	acgggacgct	ctacatccgc	aacctcgcgc	ccaaggacag	cgggcgctat	6360
gagtgcgtgg	ccgccaaact	ggtaggctcc	gcgcgcagga	cgggtgcagct	gaacgtgcag	6420
cgtgcagcag	ccaacgcgcg	catcacgggc	acctccccgc	ggaggacgga	cgtcaggtac	6480
ggaggaaccc	tcaagctgga	ctgcagcgcc	tccggggacc	cctggccgcg	catcctctgg	6540
aggctgcctg	ccaagaggat	gatcgacgcg	ctcttcagtt	ttgatagcag	aatcaagggtg	6600
tttgccaatg	ggaccctggt	ggtgaaatca	gtgacggaca	aagatgccgg	agattacctg	6660
tgcgtagctc	gaaataaggt	tgggtgatgac	tacgtggtgc	tcaaagtgga	tgtggtgatg	6720
aaaccggcca	agattgaaca	caaggaggag	aacgaccaca	aagtcttcta	cgggggtgac	6780
ctgaaagtgg	actgtgtggc	caccgggctt	cccaatcccg	agatctcctg	gagcctccca	6840
gacgggagtc	tggngaactc	cttcatgcag	tcggatgaca	gcgggtggacg	caccaagcgc	6900
tatgtcgtct	tcaacaatgg	gacactctac	tttaacgaag	tggggatgag	ggaggaagga	6960
gactacacct	gctttgctga	aaatcaggct	gggaaggacg	agatgagagt	cagagtcaag	7020
gtggtgacag	cgcccgccac	catccggaac	aagacttact	tggcggttca	ggtgccctat	7080
ggagacgtgg	tactgttagc	ctgtgaggcc	aaaggagaac	ccatgcccac	ggtgacttgg	7140
ttgtcccca	ccaacaaggt	gatccccacc	tctctgaga	agtatcagat	ataccaagat	7200
ggcactctcc	ttattcagaa	agcccagcgt	tctgacagcg	gcaactacac	ctgcctggtc	7260
aggaacagcg	cgggagagga	taggaagacg	gtgtggatc	acgtcaacgt	ccagccaccc	7320
aagatcaacg	gtaaccccaa	ccccatcacc	accgtgcggg	agatagcagc	cgggggcagt	7380
cggaaaactga	ttgactgcaa	agctgaaggc	atccccaccc	cgagggtggt	atgggctttt	7440
cccgaggggtg	tggttctgcc	agctccatac	tatggaaaac	ggatcactgt	ccatggcaac	7500
ggttcccttg	acatcaggag	tttgaggaag	agcgactccg	tccagctggt	atgcattggca	7560
cgcacagagg	gaggggaggc	gaggttgatc	gtgcagctca	ctgtcctgga	gcccattggag	7620
aaacccactct	tccacgacct	gatcagcgag	aagatcacgg	ccatggcggg	ccacaccatc	7680
agcctcaact	gctctgcgcg	ggggaccctg	acaccagccc	tgggtgtgggt	ccttcccaat	7740
ggcaccgatc	tgcagatgtg	acagcagctg	cagcgcttct	accacaaggc	tgacggcatg	7800
ctacacatta	gcggtctctc	ctcggtggac	gctggggcct	accgctgcgt	ggcccgcaat	7860
gccgctggcc	acacggagag	gctggtctcc	ctgaagggtg	gactgaagcc	agaagcaaac	7920
aagcagtatc	ataacctggt	cagcatcatc	aatggtgaga	ccctgaagct	cccctgcacc	7980
cctcccgggg	ctgggcaggg	acgtttctcc	tggacgctcc	ccaatggcat	gcatctggag	8040
ggcccccaaa	ccctgggacg	cgtttctctt	ctggacaatg	gcacctcac	ggttcgtgag	8100
gcctcggtgt	ttgacagggg	tacctatgta	tgcaggatgg	agacggagta	cggcccttcg	8160
gtcaccagca	tccccgtgat	tgtgategcc	tatcctcccc	ggatcaccag	cgcgcccacc	8220
ccggtcatct	acacccggcc	cgggaacacc	gtgaaactga	actgcatggc	tatggggatt	8280

```
<210> 2
<211> 2828
<212> PRT
<213> Human
```

<400>	2															
Met	Pro	Lys	Arg	Ala	His	Trp	Gly	Ala	Leu	Ser	Val	Val	Leu	Ile	Leu	
1				5					10					15		
Leu	Trp	Gly	His	Pro	Arg	Val	Ala	Leu	Ala	Cys	Pro	His	Pro	Cys	Ala	
			20					25					30			
Cys	Tyr	Val	Pro	Ser	Glu	Val	His	Cys	Thr	Phe	Arg	Ser	Leu	Ala	Ser	
		35					40					45				
Val	Pro	Ala	Gly	Ile	Ala	Arg	His	Val	Glu	Arg	Ile	Asn	Leu	Gly	Phe	
	50					55					60					
Asn	Ser	Ile	Gln	Ala	Leu	Ser	Glu	Thr	Ser	Phe	Ala	Gly	Leu	Thr	Lys	
65				70						75					80	
Leu	Glu	Leu	Leu	Met	Ile	His	Gly	Asn	Glu	Ile	Pro	Ser	Ile	Pro	Asp	
			85					90						95		
Gly	Ala	Leu	Arg	Asp	Leu	Ser	Ser	Leu	Gln	Val	Phe	Lys	Phe	Ser	Tyr	
			100					105					110			
Asn	Lys	Leu	Arg	Val	Ile	Thr	Gly	Gln	Thr	Leu	Gln	Gly	Leu	Ser	Asn	
		115					120					125				
Leu	Met	Arg	Leu	His	Ile	Asp	His	Asn	Lys	Ile	Glu	Phe	Ile	His	Pro	
	130					135					140					
Gln	Ala	Phe	Asn	Gly	Leu	Thr	Ser	Leu	Arg	Leu	Leu	His	Leu	Glu	Gly	
145				150						155					160	
Asn	Leu	Leu	His	Gln	Leu	His	Pro	Ser	Thr	Phe	Ser	Thr	Phe	Thr	Phe	

					165					170				175	
Leu	Asp	Tyr	Phe	Arg	Leu	Ser	Thr	Ile	Arg	His	Leu	Tyr	Leu	Ala	Glu
			180					185					190		
Asn	Met	Val	Arg	Thr	Leu	Pro	Ala	Ser	Met	Leu	Arg	Asn	Met	Pro	Leu
		195					200					205			
Leu	Glu	Asn	Leu	Tyr	Leu	Gln	Gly	Asn	Pro	Trp	Thr	Cys	Asp	Cys	Glu
	210					215					220				
Met	Arg	Trp	Phe	Leu	Glu	Trp	Asp	Ala	Lys	Ser	Arg	Gly	Ile	Leu	Lys
225					230					235					240
Cys	Lys	Lys	Asp	Lys	Ala	Tyr	Glu	Gly	Gly	Gln	Leu	Cys	Ala	Met	Cys
			245						250					255	
Phe	Ser	Pro	Lys	Lys	Leu	Tyr	Lys	His	Glu	Ile	His	Lys	Leu	Lys	Asp
			260					265					270		
Met	Thr	Cys	Leu	Lys	Pro	Ser	Ile	Glu	Ser	Pro	Leu	Arg	Gln	Asn	Arg
		275					280					285			
Ser	Arg	Ser	Ile	Glu	Glu	Glu	Gln	Glu	Gln	Glu	Glu	Asp	Gly	Gly	Ser
	290					295					300				
Gln	Leu	Ile	Leu	Glu	Lys	Phe	Gln	Leu	Pro	Gln	Trp	Ser	Ile	Ser	Leu
305					310					315					320
Asn	Met	Thr	Asp	Glu	His	Gly	Asn	Met	Val	Asn	Leu	Val	Cys	Asp	Ile
			325						330					335	
Lys	Lys	Pro	Met	Asp	Val	Tyr	Lys	Ile	His	Leu	Asn	Gln	Thr	Asp	Pro
			340					345					350		
Pro	Asp	Ile	Asp	Ile	Asn	Ala	Thr	Val	Ala	Leu	Asp	Phe	Glu	Cys	Pro
		355					360					365			
Met	Thr	Arg	Glu	Asn	Tyr	Glu	Lys	Leu	Trp	Lys	Leu	Ile	Ala	Tyr	Tyr
	370					375					380				
Ser	Glu	Val	Pro	Val	Lys	Leu	His	Arg	Glu	Leu	Met	Leu	Ser	Lys	Asp
385					390					395					400
Pro	Arg	Val	Ser	Tyr	Gln	Tyr	Arg	Gln	Asp	Ala	Asp	Glu	Glu	Ala	Leu
			405						410					415	
Tyr	Tyr	Thr	Gly	Val	Arg	Ala	Gln	Ile	Leu	Ala	Glu	Pro	Glu	Trp	Val
			420					425					430		
Met	Gln	Pro	Ser	Ile	Asp	Ile	Gln	Leu	Asn	Arg	Arg	Gln	Ser	Thr	Ala
		435					440					445			
Lys	Lys	Val	Leu	Leu	Ser	Tyr	Tyr	Thr	Gln	Tyr	Ser	Gln	Thr	Ile	Ser
	450					455					460				
Thr	Lys	Asp	Thr	Arg	Gln	Ala	Arg	Gly	Arg	Ser	Trp	Val	Met	Ile	Glu
465					470					475					480
Pro	Ser	Gly	Ala	Val	Gln	Arg	Asp	Gln	Thr	Val	Leu	Glu	Gly	Gly	Pro
			485						490					495	
Cys	Gln	Leu	Ser	Cys	Asn	Val	Lys	Ala	Ser	Glu	Ser	Pro	Ser	Ile	Phe
			500					505					510		
Trp	Val	Leu	Pro	Asp	Gly	Ser	Ile	Leu	Lys	Ala	Pro	Met	Asp	Asp	Pro
		515	</												

Glu	Ser	Val	Thr	Leu	Pro	Cys	Asn	Ala	Leu	Ala	Ile	Pro	Glu	Ala	His
		595					600					605			
Leu	Ser	Trp	Ile	Leu	Pro	Asn	Arg	Arg	Ile	Ile	Asn	Asp	Leu	Ala	Asn
		610				615					620				
Thr	Ser	His	Val	Tyr	Met	Leu	Pro	Asn	Gly	Thr	Leu	Ser	Ile	Pro	Lys
625					630					635					640
Val	Gln	Val	Ser	Asp	Ser	Gly	Tyr	Tyr	Arg	Cys	Val	Ala	Val	Asn	Gln
				645					650					655	
Gln	Gly	Ala	Asp	His	Phe	Thr	Val	Gly	Ile	Thr	Val	Thr	Lys	Lys	Gly
			660					665					670		
Ser	Gly	Leu	Pro	Ser	Lys	Arg	Gly	Arg	Arg	Pro	Gly	Ala	Lys	Ala	Leu
		675					680					685			
Ser	Arg	Val	Arg	Glu	Asp	Ile	Val	Glu	Asp	Glu	Gly	Gly	Ser	Gly	Met
		690				695					700				
Gly	Asp	Glu	Glu	Asn	Thr	Ser	Arg	Arg	Leu	Leu	His	Pro	Lys	Asp	Gln
705					710					715					720
Glu	Val	Phe	Leu	Lys	Thr	Lys	Asp	Asp	Ala	Ile	Asn	Gly	Asp	Lys	Lys
				725					730					735	
Ala	Lys	Lys	Gly	Arg	Arg	Lys	Leu	Lys	Leu	Trp	Lys	His	Ser	Glu	Lys
			740					745					750		
Glu	Pro	Glu	Thr	Asn	Val	Ala	Glu	Gly	Arg	Arg	Val	Phe	Glu	Ser	Arg
		755					760					765			
Arg	Arg	Ile	Asn	Met	Ala	Asn	Lys	Gln	Ile	Asn	Pro	Glu	Arg	Trp	Ala
		770				775					780				
Asp	Ile	Leu	Ala	Lys	Val	Arg	Gly	Lys	Asn	Leu	Pro	Lys	Gly	Thr	Glu
785					790					795					800
Val	Pro	Pro	Leu	Ile	Lys	Thr	Thr	Ser	Pro	Pro	Ser	Leu	Ser	Leu	Glu
				805					810					815	
Val	Thr	Pro	Pro	Phe	Pro	Ala	Val	Ser	Pro	Pro	Ser	Ala	Ser	Pro	Val
		820						825					830		
Gln	Thr	Val	Thr	Ser	Ala	Glu	Glu	Ser	Ser	Ala	Asp	Val	Pro	Leu	Leu
		835					840					845			
Gly	Glu	Glu	Glu	His	Val	Leu	Gly	Thr	Ile	Ser	Ser	Ala	Ser	Met	Gly
		850				855					860				
Leu	Glu	His	Asn	His	Asn	Gly	Val	Ile	Leu	Val	Glu	Pro	Glu	Val	Thr
865					870					875					880
Ser	Thr	Pro	Leu	Glu	Glu	Val	Val	Asp	Asp	Leu	Ser	Glu	Lys	Thr	Glu
				885					890					895	
Glu	Ile	Thr	Ser	Thr	Glu	Gly	Asp	Leu	Lys	Gly	Thr	Ala	Ala	Pro	Thr
		900						905					910		
Leu	Ile	Ser	Glu	Pro	Tyr	Glu	Pro	Ser	Pro	Thr	Leu	His	Thr	Leu	Asp
		915					920					925			
Thr	Val	Tyr	Glu	Lys	Pro	Thr	His	Glu	Glu	Thr	Ala	Thr	Glu	Gly	Trp
		930				935					940				
Ser	Ala	Ala	Asp												

1010					1015					1020					
Glu	Pro	Gly	Val	Pro	Gly	Gln	Ser	His	Leu	Gln	Gly	Leu	Thr	Asp	Asn
1025					1030					1035					1040
Ile	His	Leu	Val	Lys	Ser	Ser	Leu	Ser	Thr	Gln	Asp	Thr	Leu	Leu	Ile
				1045					1050					1055	
Lys	Lys	Gly	Met	Lys	Glu	Met	Ser	Gln	Thr	Leu	Gln	Gly	Gly	Asn	Met
			1060					1065					1070		
Leu	Glu	Gly	Asp	Pro	Thr	His	Ser	Arg	Ser	Ser	Glu	Ser	Glu	Gly	Gln
		1075					1080				1085				
Glu	Ser	Lys	Ser	Ile	Thr	Leu	Pro	Asp	Ser	Thr	Leu	Gly	Ile	Met	Ser
1090						1095					1100				
Ser	Met	Ser	Pro	Val	Lys	Lys	Pro	Ala	Glu	Thr	Thr	Val	Gly	Thr	Leu
1105				1110						1115					1120
Leu	Asp	Lys	Asp	Thr	Thr	Thr	Val	Thr	Thr	Thr	Pro	Arg	Gln	Lys	Val
				1125				1130						1135	
Ala	Pro	Ser	Ser	Thr	Met	Ser	Thr	His	Pro	Ser	Arg	Arg	Arg	Pro	Asn
			1140					1145					1150		
Gly	Arg	Arg	Arg	Leu	Arg	Pro	Asn	Lys	Phe	Arg	His	Arg	His	Lys	Gln
	1155					1160					1165				
Thr	Pro	Pro	Thr	Thr	Phe	Ala	Pro	Ser	Glu	Thr	Phe	Ser	Thr	Gln	Pro
1170						1175					1180				
Thr	Gln	Ala	Pro	Asp	Ile	Lys	Ile	Ser	Ser	Gln	Val	Glu	Ser	Ser	Leu
1185				1190						1195					1200
Val	Pro	Thr	Ala	Trp	Val	Asp	Asn	Thr	Val	Asn	Thr	Pro	Lys	Gln	Leu
				1205				1210						1215	
Glu	Met	Glu	Lys	Asn	Ala	Glu	Pro	Thr	Ser	Lys	Gly	Thr	Pro	Arg	Arg
			1220					1225					1230		
Lys	His	Gly	Lys	Arg	Pro	Asn	Lys	His	Arg	Tyr	Thr	Pro	Ser	Thr	Val
	1235					1240					1245				
Ser	Ser	Arg	Ala	Ser	Gly	Ser	Lys	Pro	Ser	Pro	Ser	Pro	Glu	Asn	Lys
	1250					1255				1260					
His	Arg	Asn	Ile	Val	Thr	Pro	Ser	Ser	Glu	Thr	Ile	Leu	Leu	Pro	Arg
1265				1270						1275					1280
Thr	Val	Ser	Leu	Lys	Thr	Glu	Gly	Pro	Tyr	Asp	Ser	Leu	Asp	Tyr	Met
				1285				1290						1295	
Thr	Thr	Thr	Arg	Lys	Ile	Tyr	Ser	Ser	Tyr	Pro	Lys	Val	Gln	Glu	Thr
			1300					1305					1310		
Leu	Pro	Val	Thr	Tyr	Lys	Pro	Thr	Ser	Asp	Gly	Lys	Glu	Ile	Lys	Asp
		1315				1320					1325				
Asp	Val	Ala	Thr	Asn	Val	Asp	Lys	His	Lys	Ser	Asp	Ile	Leu	Val	Thr
	1330					1335				1340					
Gly	Glu	Ser	Ile	Thr	Asn	Ala	Ile	Pro	Thr	Ser	Arg	Ser	Leu	Val	Ser
1345				1350						1355					1360
Thr	Met	Gly	Glu	Phe	Lys	Glu	Glu	Ser	Ser	Pro	Val	Gly	Phe	Pro	Gly
			1365												

Thr	Thr	Leu	Ser	Ile	Lys	Val	Glu	Val	Ala	Ser	Ser	Gln	Ala	Glu	
				1445				1450					1455		
Thr	Thr	Thr	Leu	Asp	Gln	Asp	His	Leu	Glu	Thr	Thr	Val	Ala	Ile	Leu
			1460					1465					1470		
Leu	Ser	Glu	Thr	Arg	Pro	Gln	Asn	His	Thr	Pro	Thr	Ala	Ala	Arg	Met
			1475					1480					1485		
Lys	Glu	Pro	Ala	Ser	Ser	Ser	Pro	Ser	Thr	Ile	Leu	Met	Ser	Leu	Gly
			1490					1495				1500			
Gln	Thr	Thr	Thr	Thr	Lys	Pro	Ala	Leu	Pro	Ser	Pro	Arg	Ile	Ser	Gln
1505					1510						1515				1520
Ala	Ser	Arg	Asp	Ser	Lys	Glu	Asn	Val	Phe	Leu	Asn	Tyr	Val	Gly	Asn
				1525					1530					1535	
Pro	Glu	Thr	Glu	Ala	Thr	Pro	Val	Asn	Asn	Glu	Gly	Thr	Gln	His	Met
			1540					1545					1550		
Ser	Gly	Pro	Asn	Glu	Leu	Ser	Thr	Pro	Ser	Ser	Asp	Arg	Asp	Ala	Phe
			1555					1560				1565			
Asn	Leu	Ser	Thr	Lys	Leu	Glu	Leu	Glu	Lys	Gln	Val	Phe	Gly	Ser	Arg
			1570					1575				1580			
Ser	Leu	Pro	Arg	Gly	Pro	Asp	Ser	Gln	Arg	Gln	Asp	Gly	Arg	Val	His
1585					1590					1595					1600
Ala	Ser	His	Gln	Leu	Thr	Arg	Val	Pro	Ala	Lys	Pro	Ile	Leu	Pro	Thr
				1605					1610					1615	
Ala	Thr	Val	Arg	Leu	Pro	Glu	Met	Ser	Thr	Gln	Ser	Ala	Ser	Arg	Tyr
			1620					1625				1630			
Phe	Val	Thr	Ser	Gln	Ser	Pro	Arg	His	Trp	Thr	Asn	Lys	Pro	Glu	Ile
			1635					1640				1645			
Thr	Thr	Tyr	Pro	Ser	Gly	Ala	Leu	Pro	Glu	Asn	Lys	Gln	Phe	Thr	Thr
		1650				1655					1660				
Pro	Arg	Leu	Ser	Ser	Thr	Thr	Ile	Pro	Leu	Pro	Leu	His	Met	Ser	Lys
1665					1670					1675					1680
Pro	Ser	Ile	Pro	Ser	Lys	Phe	Thr	Asp	Arg	Arg	Thr	Asp	Gln	Phe	Asn
				1685					1690					1695	
Gly	Tyr	Ser	Lys	Val	Phe	Gly	Asn	Asn	Asn	Ile	Pro	Glu	Ala	Arg	Asn
			1700					1705					1710		
Pro	Val	Gly	Lys	Pro	Pro	Ser	Pro	Arg	Ile	Pro	His	Tyr	Ser	Asn	Gly
		1715						1720				1725			
Arg	Leu	Pro	Phe	Phe	Thr	Asn	Lys	Thr	Leu	Ser	Phe	Pro	Gln	Leu	Gly
		1730				1735					1740				
Val	Thr	Arg	Arg	Pro	Gln	Ile	Pro	Thr	Ser	Pro	Ala	Pro	Val	Met	Arg
1745					1750					1755					1760
Glu	Arg	Lys	Val	Ile	Pro	Gly	Ser	Tyr	Asn	Arg	Ile	His	Ser	His	Ser
				1765					1770					1775	
Thr	Phe	His	Leu	Asp	Phe	Gly	Pro	Pro	Ala	Pro	Pro	Leu	Leu	His	Thr
			1780					1785							

[illegible]

Met	Gln	Ser	Asp	Asp	Ser	Gly	Gly	Arg	Thr	Lys	Arg	Tyr	Val	Val	Phe
2290						2295				2300					
Asn	Asn	Gly	Thr	Leu	Tyr	Phe	Asn	Glu	Val	Gly	Met	Arg	Glu	Glu	Gly
2305					2310					2315					2320
Asp	Tyr	Thr	Cys	Phe	Ala	Glu	Asn	Gln	Val	Gly	Lys	Asp	Glu	Met	Arg
				2325					2330						2335
Val	Arg	Val	Lys	Val	Val	Thr	Ala	Pro	Ala	Thr	Ile	Arg	Asn	Lys	Thr
			2340					2345					2350		
Tyr	Leu	Ala	Val	Gln	Val	Pro	Tyr	Gly	Asp	Val	Val	Thr	Val	Ala	Cys
		2355					2360					2365			
Glu	Ala	Lys	Gly	Glu	Pro	Met	Pro	Lys	Val	Thr	Trp	Leu	Ser	Pro	Thr
2370						2375					2380				
Asn	Lys	Val	Ile	Pro	Thr	Ser	Ser	Glu	Lys	Tyr	Gln	Ile	Tyr	Gln	Asp
2385					2390					2395					2400
Gly	Thr	Leu	Leu	Ile	Gln	Lys	Ala	Gln	Arg	Ser	Asp	Ser	Gly	Asn	Tyr
				2405					2410						2415
Thr	Cys	Leu	Val	Arg	Asn	Ser	Ala	Gly	Glu	Asp	Arg	Lys	Thr	Val	Trp
			2420					2425					2430		
Ile	His	Val	Asn	Val	Gln	Pro	Pro	Lys	Ile	Asn	Gly	Asn	Pro	Asn	Pro
		2435					2440					2445			
Ile	Thr	Thr	Val	Arg	Glu	Ile	Ala	Ala	Gly	Gly	Ser	Arg	Lys	Leu	Ile
		2450				2455					2460				
Asp	Cys	Lys	Ala	Glu	Gly	Ile	Pro	Thr	Pro	Arg	Val	Leu	Trp	Ala	Phe
2465					2470					2475					2480
Pro	Glu	Gly	Val	Val	Leu	Pro	Ala	Pro	Tyr	Tyr	Gly	Asn	Arg	Ile	Thr
				2485					2490						2495
Val	His	Gly	Asn	Gly	Ser	Leu	Asp	Ile	Arg	Ser	Leu	Arg	Lys	Ser	Asp
			2500					2505					2510		
Ser	Val	Gln	Leu	Val	Cys	Met	Ala	Arg	Asn	Glu	Gly	Gly	Glu	Ala	Arg
		2515					2520					2525			
Leu	Ile	Val	Gln	Leu	Thr	Val	Leu	Glu	Pro	Met	Glu	Lys	Pro	Ile	Phe
		2530				2535					2540				
His	Asp	Pro	Ile	Ser	Glu	Lys	Ile	Thr	Ala	Met	Ala	Gly	His	Thr	Ile
2545					2550					2555					2560
Ser	Leu	Asn	Cys	Ser	Ala	Ala	Gly	Thr	Pro	Thr	Pro	Ser	Leu	Val	Trp
				2565					2570						2575
Val	Leu	Pro	Asn	Gly	Thr	Asp	Leu	Gln	Ser	Gly	Gln	Gln	Leu	Gln	Arg
			2580					2585					2590		
Phe	Tyr	His	Lys	Ala	Asp	Gly	Met	Leu	His	Ile	Ser	Gly	Leu	Ser	Ser
		2595				2600						2605			
Val	Asp	Ala	Gly	Ala	Tyr	Arg	Cys	Val	Ala	Arg	Asn	Ala	Ala	Gly	His
					2615						2620				
Thr	Glu	Arg	Leu	Val	Ser	Leu	Lys	Val	Gly	Leu	Lys	Pro	Glu	Ala	Asn
2625					2630										

```
<210> 3
<211> 27
<212> DNA
<213> Artificial Sequence
```

```
<400> 3
qqagtcctgt qgcattccacg aaactac
```

```
<210> 4
<211> 25
<212> DNA
<213> Artificial Sequence
```

```
<400> 4
cacatctgct qqaaggtgga cagcg
```

```
<210> 5
<211> 26
<212> DNA
<213> Artificial Sequence
```

```
<400> 5
catgggcaca aactgactca tggctg
```

```
<210> 6
<211> 28
<212> DNA
<213> Artificial Sequence
```

<223> PCR primer

gagaggagag gaaggagaaa actgcatc

28

<211> 37

<212> DNA

<213> Artificial Sequence

<223> PCR primer

ttgcggccgc gccaccatgc ccaagcgcgc gcactgg

37

<211> 28

<212> DNA

<213> Artificial Sequence

<223> PCR primer

tcaatactcc tgctcctggt ctgtctca

28